



## METHODS AND REAGENTS FOR REGULATION OF CELLULAR RESPONSES IN BIOLOGICAL SYSTEMS

### STATEMENT REGARDING U.S. GOVERNMENT FUNDING

5

This invention was made at least in part with Funding from the United States government through National Institute of Health grant GM55984. The United States government has certain rights in this invention.

### CROSS-REFERENCE TO RELATED APPLICATIONS

This application takes priority under 35 U.S.C. 119(d) from U.S. provisional application serial number 60/191,014 filed March 21, 2000 which is incorporated by reference herein to the extent that it is not inconsistent with the disclosure herein.

### BACKGROUND OF THE INVENTION

15  
20  
25

A variety of biological processes are mediated by the binding of one chemical or biological species, macromolecule or particle (e.g., a cell, virus or virion) to another chemical or biological species, macromolecule or particle. In many cases there is evidence that the valency of the binding may be an important aspect of the mechanism of the mediation of the biological process. The present invention relates to compounds and methods for selectively varying the valency such interactions employing multivalent ligands to which a plurality of chemical or biological species involved in binding to other chemical or biological species (generally designated recognition elements, RE, herein) attached in a controlled fashion, with control over the number of RE, the spacing of RE and the relative orientation of RE. Certain recognition elements are involved directly or indirectly in biological signaling processes other recognition elements are involved simply in facilitating binding that is associated with the biological process. This invention then is generally related to the control of biological processes by controlling the structure of such multivalent ligands. The multivalent ligands of this invention have applications particularly in cell signaling processes and more generally in macromolecular assembly of recognition elements that are involved in biological processes.

Cells need to continuously sense and respond to changes in their environment. For this purpose, cells use a multitude of cell surface, transmembrane and cytoplasmic receptors. These receptors typically recognize proteins, peptides, saccharides, nucleic acids, or other small molecules but, in some cases, receptors may also recognize changes in redox potential, temperature, and osmolarity. The binding of a ligand to these receptors results in changes in the activity of the cell such as migration, activation, metabolism, protein production, differentiation, proliferation, and cell death. This is a central paradigm of cell biology and these cellular responses allow the cell (or the multicellular organism) to properly respond to environmental changes.

The mechanisms by which ligands promote cellular processes are of great interest to elucidate their roles in the regulation of cellular responses. One common way in which these systems are regulated is by the spatial organization of the receptors. Ligand binding can change the relative orientations and/or conformations of the cell surface receptors, activating a response. Biological responses ranging from immune recognition, cell adhesion and migration, and proliferation, among others, rely on the reorientation or change in distribution (e.g., localization) of cell surface receptors that occurs upon ligand binding. Ligand reorientation can be the event that transmits signals and facilitates the cellular response.

A common example of this is found in the growth factor receptors, which govern cell proliferation. Certain divalent growth factors, such as erythropoietin (EPO), bind to two cell surface growth factor receptors (EPOR) simultaneously and bring those receptors into proximity. This ligand reorganization triggers a signal transduction cascade that involves cross-phosphorylation of the receptors in the dimer. In the EPO example a ligand, which is only capable of binding to one receptor, is incapable of eliciting the response.

A particularly interesting feature of some ligands is valency, which herein refers generally to an interplay between the net number of recognition sites in a ligand for binding to receptors (e.g., epitopes) and the density and spacing of those sites in the ligand. Ligands often possess multiple receptor binding sites. This allows multivalent interactions between the ligand and multiple receptors which may determine the kind and intensity of biological response to that ligand. Often in these systems, monovalent ligands lack any biological activity. Researchers have explored ligands which vary in valency, at least in the sense of increasing the number of recognition sites. Typically the ligands examined have been either small, low valency compounds, such as antibodies or dimerizing agents, or large

heterogeneous compounds, such as protein conjugates, polymers, or functionalized surfaces. Work with defined low valency compounds has led to the realization of the extent of regulation by changes in receptor orientation and work with large undefined multivalent ligands has indicated that increasing the net number of recognition sites (e.g., epitopes) can often result in increased effects in many systems.

Cells require fine control over their cellular processes in order to avoid over- or under-stimulation. In the immune response, for example, immune cell function must be closely regulated to avoid unfavorable autoreactivity or clonal anergy. Cells utilize features of the interaction of receptors with ligands to regulate their responses. For example, increased synthetic ligand density has been shown to more effectively activate the response of certain cells to the ligand. Nature may utilize ligand valency to control biological responses in a defined manner. Thus, selective control of biological responses may be achieved through control of ligand valency. Previously described multivalent ligands have, however, not allowed exploration of this fine-tuned control in biological systems.

This invention provides for the generation of synthetic ligands with distinct valencies and controlled features which can be used to systematically alter and/or control biological responses initiated or triggered by binding to cell surface receptors. In particular, the synthetic ligands of this invention allow for access to the finer control exhibited by natural ligands. Access to these features in a synthetic ligand not only expands our understanding of the natural function of these systems, but also leads to selectively designed effector molecules (multivalent ligands) for use in therapeutic and non-therapeutic applications that take advantage of the ability to regulate a wide variety of biological responses.

### SUMMARY OF THE INVENTION

This invention provides multivalent ligands which carry or display at least one recognition element (RE), and preferably a plurality of recognition elements, for binding directly or indirectly to cells or other biological particles or more generally by binding to any biological molecule. The multivalent ligands provided can most generally function for binding or targeting to any biological particle or molecule and particularly to targeting of cells or cell types or viruses, for cell aggregation and generally for macromolecular assembly of biological macromolecules. The multivalent ligands of this invention are generally applicable for creating scaffolds (assemblies) of chemical or biological species, including

without limitation, antigens, epitopes, ligand binding groups, ligands for cell receptors (cell surface receptors, transmembrane receptors and cytoplasmic receptors), various macromolecules (nucleic acids, carbohydrates, saccharides, proteins, peptides, etc.). In these scaffolds, the number, spacing, relative positioning and relative orientation of recognition elements can be controlled.

In a more specific embodiment, multivalent ligands are provided which carry or display at least one signal recognition element (SRE), and preferably a plurality of signal recognition elements, and modulate biological responses in biological systems. Signal recognition elements provide for binding to a cell surface receptor and alone or in combination with other SRE affect a biological response in a biological system. SRE include chemical or biochemical species recognized as signals by a cell, i.e., through binding one or more cell receptors, particularly one or more cell surface receptors. These multivalent ligands can act generally as effectors of biological responses in biological systems. The multivalent ligands provided can function to activate, initiate or trigger a biological response, to inhibit a response, to enhance or attenuate a response, or to change the nature of a response. A multivalent ligand of this invention can also affect a response mediated through a cell surface receptor to which it does not itself bind.

The invention provides methods for labeling or targeting of cells with functional elements (FE). The invention also provides methods for inducing or enhancing cell aggregation or alternatively for inhibiting or preventing cell aggregation. The invention further provides methods for inducing, modulating and/or regulating biological responses in biological systems. Each of these methods employ multivalent ligands. Preferred multivalent ligands of this invention have defined or controlled valency, in which structural features of the ligand are selected or controlled, including the number, density, spacing and orientation of recognition elements (RE and SRE) for binding to receptors, to simply bind to a cell or to obtain a desired type of biological response or level of response.

Scaffolded multivalent ligands of this invention which comprise a plurality of RE, SRE or both optionally in combination with FE can be employed in a variety of diagnostic and clinical applications, in particular in blood typing and in pathogen detection. The multivalent ligands herein can be employed in the detection of various biological molecules and particles (cells and viruses) and in a variety of assay methods (histology, Western blots,

PCR assays, ELISA assays, agglutination assays, among others). In general, increased in valency in such ligands will be associated with an increase in assay or diagnostic sensitivity.

Multivalent ligands comprise one or more structural or functional groups which act as recognition elements (RE) for binding to cell surface receptors, optionally in combination with one or more signal recognition elements (SRE), or one or more functional elements (FE), or both. SREs are a subset of REs that, alone or in combination with other SREs (REs or FEs) in a multivalent ligand, can induce intracellular and/or intercellular biological responses. Multivalent ligands of this invention carrying one or more SRE (optionally in combination with one or more RE, one or more different SRE or one or more FE) can initiate a biological response in a cell. Alternatively, these multivalent ligands can modulate the response of a cell in the presence of one or more natural chemical or biochemical signals, for example, by enhancing, decreasing or inhibiting the response. In specific embodiments, multivalent ligands of this invention are designed to change the level or type of response that is induced in a cell by a selected chemical or biochemical signal.

Multivalent ligands of this invention most generally comprise a molecular scaffold to which a plurality of REs, SREs or both (optionally in combination with FEs) are bonded either by covalent or non-covalent interactions. The number, density and spacing of the RE, SRE and FE on the scaffold can be controlled, typically by selective synthesis of desired ligands. The molecular scaffold can be linear, branched or cyclic providing different geometries of presentation of RE and/or SREs to cells. In preferred embodiments, molecular scaffolds are polymers comprising a plurality of monomers. Molecular scaffold of the multivalent ligands of this invention include polymers in which all of the monomers are the same or copolymers containing a mixture of different monomers. Molecular scaffolds can also include block copolymers in which different regions (or portions) of the scaffold are composed of different monomers. Molecular scaffolds prepared by ROMP methods, as illustrated in several formulas herein, are preferred.

Molecular scaffolds can be hydrophobic or can be made to be more hydrophilic by substitution (particularly of the polymer backbone) with polar substituents, such as -OH. The scaffold can be substituted, in general, with any groups that do not interfere with RE or SRE activity, e.g. binding to a receptor. Substitution of the scaffold can be controlled to adjust the physical properties, e.g., solubility, of the multivalent ligand. REs, SREs and FEs may be directly attached to a scaffold or attached to the scaffold via linker groups. The linker group

provides functional groups for bonding to the scaffold and for bonding to REs, SREs and/or FEs and can also affect solubility of the multivalent ligand. The linker can also provide a defined spacer to minimize undesired interactions among REs, SREs or FEs or between the attached elements and the scaffold or to provide structural flexibility with respect to orientation of attached elements.

In specific embodiments, the molecular scaffold comprises a plurality of repeated units (monomers) to each of which an RE or SRE is bonded. In general, the molecular scaffold functions to hold the signals in proximity to each other and does not interact directly in the modulation of the biological response. However, physical (e.g., solubility) or chemical (e.g., stability) properties of the multivalent ligands can be varied by selection of the structure of the scaffold or by introducing substituents (e.g., polar, non-polar) along the scaffold.

In one embodiment, the multivalent ligands have only one type of RE or SRE in the ligand. These multivalent ligands include dimers, trimers, tetramers and polymers (including relatively short oligomers having 5 or more monomers) or longer polymers having 25, 50, 100 or more monomers. Preferred multivalent ligands carrying one type of RE or SRE carry about 10 or more of such REs or SREs. In this embodiment, the repeating units (or monomers) of the multivalent ligand are preferably the same.

In another embodiment, the invention provides multivalent ligands that carry more than one type of RE, more than one type of SRE or a combination of RE and SRE. These multivalent ligands also include dimers (carrying one of each RE or SRE or an RE and an SRE), trimers, tetramers and block polymers (including relatively short oligomers having 5 monomers or more) or longer polymers having 25, 50, 100 or more monomers. These multivalent ligands may also have spacer regions (with monomers that do not carry any RE or SRE group) along the scaffold to separate regions carrying a first RE or SRE from regions carrying a second RE or SRE. Monomers in spacer regions may carry a functional element (FE), may be unsubstituted or may carry a non-reactive, non-functional group. A given multivalent ligand can generally contain any number of different REs, SREs, or both, however those carrying 2 or 3 different RE or SRE are of most interest.

In other embodiments, the invention provides multivalent ligands that carry one or more RE or SRE, which may be the same or different, but also carry functional elements other than RE or SRE. These functional elements (FE) can, for example, exhibit a variety of chemical or biochemical functions (different from those of REs or SREs). They can, for example, provide one or more fluorescent or radiolabels, provide one or more groups with

latent reactive groups, or provide one or more enzymatic functions. Substitution of monomers with FEs can also provide for spacing of SREs.

Recognition elements (RE) are any chemical or biological species (e.g., molecules or portions thereof) that alone or in combination with one or more other REs, recognize and bind to a cell surface receptor. RE can, for example, include all or a portion of a ligand active for binding to a cell surface receptor. Signal recognition elements (SRE) are any chemical or biochemical species that, alone or in combination with one or more other SREs, induce a biological response in or from a cell and include biological molecules (proteins, glycoproteins, peptides, amino acids, nucleic acids, saccharides, cytokines, growth factors, hormones, and various derivatives thereof) and which may be portions of larger biological species (protein fragments, epitopes, antigenic determinant, etc.) and various chemical species (haptens, naturally-occurring drugs, synthetic drugs) and species that act as functional mimics of biological molecules (e.g., peptoids, phosphorothioates). SRE are typically RE which bind to a cell surface receptor, but in contrast to RE, SRE affect a biological response in the cell.

Multivalent ligands of this invention can function to reorganize and/or cluster cell receptors. In this regard the RE or SRE on the multivalent ligand will be a ligand of the cell receptor. In certain cases, clustering or reorganization of receptors modulates the cell's response to a given SRE. Clustering or reorganization of receptors by a multivalent ligand of this invention can also modulate the response of a cell to another signal or another ligand. Through clustering or other structural reorientation or reorganization of cell surface receptors, a multivalent ligand of this invention can enhance or inhibit the cell's response to another signal or ligand. For example, multivalent ligands of this invention that function as chemoattractants can enhance the response of a cell to another chemoattractant.

A given cell receptor may mediate more than one biological response. The multivalent ligands of this invention that carry ligands which bind to given cell receptor, but which do not induce a biological response mediated by that receptor may be employed to inhibit the biological response.

Multivalent ligands that carry more than one type of SRE can be used to simultaneously or sequentially induce more than one biological response in or from a cell. Alternatively, the cellular response to one SRE can be modified by the cellular response to another SRE. Multivalent ligands carrying two or more different SREs can function, for example, to reorganize different receptors on the cell surface, which can result in modulation

of cellular response to one or more SREs. Similarly, in multivalent ligands carrying FE, in addition to one or more SRE, the response to an SRE can be modified by the presence of FE.

10 Multivalent ligands of this invention can be employed in methods to modulate signal transduction processes (i.e., the transmission of information between the outside and the inside of a cell and between cells, in biological systems) in prokaryotic or eukaryotic cells. The methods can be practiced *in vivo*, *in vitro* or *ex vivo* (where cells are removed from a natural environment, including a multicellular organism, and are intended once treated to be returned to that environment). For example, chemotaxis or cell migration responses to SREs can be modulated. Such methods are applicable to both prokaryotic (e.g., Gram negative, as well as Gram positive bacteria), eukaryotic microorganisms (including, without limitation, eukaryotic parasites and pathogens of various organisms, including mammals), and eukaryotic cells of larger organisms including those of mammals, and specifically including those of humans (e.g., leukocytes, lymphocytes, endothelial cells, and epithelial cells). Multivalent ligands that modulate responses in bacterial cells or in eukaryotic cells, including eukaryotic pathogens or parasites, can be used to inhibit proliferation, colonization, migration, or biofilm formation by the bacterium, or eukaryotic pathogen or parasite and, as a consequence, can inhibit infection or colonization by such microorganisms.

20 Multivalent ligands can also be used to promote or inhibit cell differentiation, cell proliferation and/ or cell death (e.g., apoptosis). Multivalent ligands that modulate responses in eukaryotic cells of larger organisms can be used to inhibit undesired cell proliferation, undesired migration and undesired formation of cell to cell junctions or to promote or enhance desired cell proliferation, desire migration and desired formation of cell junctions dependent upon the selection of SRE and other FE in the multivalent ligand.

30 Pharmaceutical and therapeutic compositions which comprise a pharmaceutically acceptable carrier and an amount of a multivalent ligand effective for modulating cell proliferation, colonization, migration, cell to cell junction formation and/or biofilm formation by eukaryotic or prokaryotic cells are encompassed by this invention. Specific pharmaceutical or therapeutic compositions include those which comprise an amount of a multivalent ligand effective for inhibiting or disrupting undesired cell proliferation, colonization, migration, cell to cell junction formation and/or biofilm formation by eukaryotic or prokaryotic cells. Pharmaceutical compositions that retard or inhibit infections by bacteria or eukaryotic parasites or pathogens are of particular interest. Two or more



multivalent ligands of this invention can be combined in such pharmaceutical compositions to provide for combined effect and benefit.

Cell migration, adhesion and the formation of cell to cell junctions are involved in cancer growth and metastasis. Multivalent ligands that modulate such processes can be employed in methods and pharmaceutical compositions for inhibition of cancer growth and metastasis. Again such pharmaceutical compositions include those which comprise an amount of a multivalent ligand that is effective for inhibiting cancer cell growth, adhesion or migration. Two or more multivalent ligands of this invention can be combined in such pharmaceutical compositions to provide for combined effect and benefit.

Multivalent ligands of this invention can modulate immune responses in animals (including mammals and particularly in humans) by valency-dependent interaction with cells that function in the immune system (e.g., leukocytes and lymphocytes). In particular, multivalent ligands of this invention can modulate the response of leukocytes, including neutrophils, to chemoattractants (including derivatized peptides, such as N-formyl peptides, and N-acyl peptides) and can modulate the activation and deactivation of B-cells and/or T-cells. B-cell and/or T-cell activation can be performed *in vivo*, *in vitro* and/or *ex vivo*.

The invention also provides libraries of multivalent ligands in which the members of the libraries are varied, for example, in the number and/or relative positioning of RE or SRE, the presence and/or positioning of spacers, in the number of repeating units or monomers (e.g.,  $n$  or  $n+m$  in formulas below) and in the presence or number of FE. Libraries of multivalent ligands which span a range of defined sizes, numbers of repeating units or monomers, numbers of RE or SRE, combinations of RE or SRE, combinations of RE, SRE and FE and spacing of attached elements, (RE, SRE and any FE) are of particular interest. Libraries prepared using ROMP-methods are of particular interest and application. Using various selection and screening methods that are understood in the art, these libraries can be selected or screened for multivalent ligands in the library which exhibit desired modulation in a given biological system. Furthermore, the results obtained from such screens, i.e., the number of RE required for cell aggregation, the number of SRE's required for induction or inhibition, and other structure/function relationships, can be used in the design and synthesis of additional multivalent ligands.

#### BRIEF DESCRIPTION OF THE DRAWINGS

July 14/11  
Figure 1 schematically illustrates several ways in which multivalent ligand of this invention can function in macromolecular assembly and as effectors of biological responses.

Figure 2A: Results of video microscopy motion analysis experiments (I). Bacteria (*Escherichia coli*) were treated with buffer alone, galactose, or compound 1 or 3 (Scheme 1) at the indicated saccharide concentrations. The results represent the average from at least five independent experiments performed in triplicate. Error bars represent the deviation between per-second averages during the ten second interval.

Figures 2B-E: Selected sample paths for bacteria (Gram Negative, *E. coli*) treated with buffer alone (B); 1 mM galactose (C); or 1mM compound 1 (D); or 1 mM compound 3 (E). Sample paths are derived from motion of representative bacteria from a treated bacterial population.

Figures 3A and 3B: Results of *E. coli* capillary accumulation assays. The number of bacteria accumulated is plotted versus the concentration of the attractant (galactose or compounds 1 -4, Scheme 1) calculated on a saccharide residue basis. (A): Results are shown for capillaries filled with buffer alone (0), compound 1 (1), and compound 2 (2) or (B): buffer alone (0), compound 3 (3) and compound 4 (4) at the indicated concentrations. The vertical line at 1 mM indicates the concentration of maximum chemotaxis for the monomeric compound 1. The concentrations used in this assay are not directly comparable to those used in the motion analysis experiments (see Figure 2A), because the gradient formed in the capillary assay is not defined. Results are the average of 3 to 6 experiments performed in duplicate and error bars represent a single standard deviation. Partial permeabilization was required to obtain chemotaxis towards 4, and was utilized for all experiments [57].

Figure 4: Results of *B. subtilis* capillary accumulation assays using ROMP-derived glucose ligands (compound 5 -7, Scheme 1). Buffer alone, glucose, or glucose-bearing compounds 5-7 were used as attractants in the capillary accumulation assay. Results are shown for glucose (G), compound 5 (5), compound 6 (6), and compound 7 (7). Results are the average of at least four trials performed in duplicate and error bars represent single standard deviations.

Figures 5A and B: Results of video microscopy motion analysis experiments (II). (A): Bacteria (*E. coli*) were treated with increasing concentrations of serine ( $\mu\text{M}$ ) after initial treatment (followed by a 2 min adaptation period) with buffer alone (■) or 10 $\mu\text{M}$  attractant: galactose (●), compound 1 (10mer, ▲) or compound 3 (25 mer, ◆); (B) Bar graph of data for angular mean velocity taken from Fig. 6A at serine concentration 1 $\mu\text{M}$ . Initial treatment with

compound 3 results in a significant enhancement of bacterial response to serine. Angular mean velocities varied approximately 14% between experiments performed on different days.

Figure 6: Multivalent ligands bind specifically to chemoreceptors and induce receptor reorganization. The illustration schematic represents fluorescently labeled 8 (10, 590 nm emission) bound to Trg (11) via GGBP(12). Trg is labeled with anti-Tsr antibody (13, 530 nm emission).

Figures 7A-D: Model of receptor reorganization by synthetic ligands. (A) Chemoreceptors are observed to form dimers (or multimers) (20) in the plasma membrane of *E. coli* and each dimer appears to interact with a single periplasmic binding protein (21) [59, 60]. Monovalent galactose ligands, such as galactose and compound 1 (22), interact with Trg through GGBP binding, inducing signal transduction from chemoreceptor dimers; (B) Multivalent galactose compounds, such as compound 2, that cannot span the distance needed to reorganize the receptors (23) generate signals from individual dimers, as in (A); (C) Multivalent ligands of sufficient lengths (24), such as compounds 3 and 4, are able to reorganize the chemoreceptors into discrete clusters (25) at the plasma membrane; (D) Extending the valency of a multivalent ligand (26) likely increases the extent of reorganization and, therefore, the bacterial response.

Figure 8: illustrates various designs for molecular scaffolds that can be employed in the multivalent ligands of this invention. These types of scaffolds can be constructed, for example, employing alicyclic, aromatic (including heteroaromatic) ring systems and combinations thereof. Scaffolds provide the geometry of presentation of two or more REs, SREs or both. Linkers may have varying structures and, for example, be rigid, flexible or branched. In each of the illustrated structures any of a rigid, flexible or branched linker can be employed. Each branched linker may be attached to more than one RE, SRE (or FE). In each structure, one or more FE (so long as at least one RE or SRE remains) can replace one or more RE or SRE.

Figures 9A-C: Illustrate models of the ability of multivalent ligands to activate or inhibit cell aggregation in a valency- and concentration-dependent fashion; (A) Monovalent ligands (31) (such as 9) are necessarily inhibitory if they bind to ConA (30); (B) Multivalent ligands 32 (such as 12) at sufficiently low concentrations and optimal stoichiometry with ConA may allow cell aggregation (34), despite their occupation of ConA binding sites; (C) At increased concentrations of multivalent ligands 32 (approximately 5  $\mu$ M in the case of 10-12) ConA sites become saturated (35), disassembling clusters and inhibiting cell aggregation.

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995  
1000

Figure 10: Bar graph illustrating that ConA clusters assembled on ROMP-derived scaffolds are able to form aggregates of Jurkat cells. Percent of Jurkat cells present in aggregates is plotted against the treatment. ConA at 100  $\mu\text{g/mL}$  or 5  $\mu\text{g/mL}$  is able to form aggregates. Aggregate formation could be inhibited by addition of 50 mM methyl  $\alpha$ -D-mannopyranoside ( $\alpha$  man). Compounds 9-12 were added to a final mannose concentrations of 0.5  $\mu\text{M}$  or 5  $\mu\text{M}$  along with a final ConA concentration of 5  $\mu\text{g/mL}$ . Results are the average of at least three independent experiments and error bars represent single standard deviations.

Figure 11: Controlling ConA-mediated erythrocyte agglutination. A graph of macroscopic aggregation index (%MAI) as a function of time after contact with cells (sec) for treatments with ConA alone or ConA in combination with ligand compound 13 (Scheme 1, mannose containing ligand with  $n = 100$ ). The concentration of Con A used was 5  $\mu\text{g/mL}$  (53 nM, based on ConA tetramer) and ligand (530 nM, based on saccharide). Thus, the ratio of mannose (in the ligand) to ConA tetramer in the experiment was 10:1. Addition of the multivalent ligand significantly enhanced erythrocyte agglutination.

Figure 12: Enhancement of Cell Toxicity of ConA by a Multivalent Ligand. A bar graph indicating % cell viability of PC12 cells as a function of various treatments. "HBS" is the medium control;" ConA" is treatment with 0.1  $\mu\text{M}$  ConA (based on Con A tetramer) in HBS medium; "Compound 11" is treatment with 4  $\mu\text{M}$  compound 11 (concentration based on saccharide) in HBS; "ConA + Compound 11" is treatment with 0.1  $\mu\text{M}$  ConA and 4  $\mu\text{M}$  compound 11 in HBS. Addition of the multivalent ligand which binds to ConA significantly enhances ConA toxicity.

#### DETAILED DESCRIPTION OF THE INVENTION

The multivalent ligands of this invention are molecular scaffolds to which a plurality of functional or structural groups, particularly RE and/or SREs, are bonded, to present a display of the functional or structural groups in a productive manner. The scaffold can in general be formed from any chemical or biological species that provides the desired orientation of display. In addition to linear arrays, the scaffolds can be chosen to provide arrays of functional groups with selected non-linear presentation. See, for example, the various non-linear scaffold structures illustrated in Fig. 8.

The functional or structural groups may be bonded to the scaffold in a symmetric or unsymmetric array. The scaffold may comprise a relatively small organic molecule, such as an aromatic ring system (including benzene, naphthalene and fused and non-fused aromatics). Various fused aromatic systems can provide a wide range of different display orientations

with functional groups bonded at selected positions on the ring system. Alternatively, saturated ring systems (e.g., cyclohexanes), heterocycles (e.g., carbohydrates), or alicyclic compounds (e.g., tris(hydroxymethyl)aminomethane) can also be used. Molecular scaffolds more typically comprise a plurality of repeating units or monomers, e.g., are polymers or oligomers. The molecular scaffold then carries a plurality of functional or structural groups bonded to repeating units or monomers. The functional groups are bonded covalently or noncovalently to the scaffold and can comprise a plurality of recognition elements (RE), or signal recognition elements (SRE), and can optionally comprise other functional elements (FE).

10 The RE, SRE and any FE can be bonded on to the molecular scaffold randomly or to a pre-selected pattern in which the arrangement of the RE, SRE and FE along the length of the scaffold matches a selected pattern, e.g., alternating different SRE or RE, selected spacing of different SRE or RE and the like).

20 The molecular scaffold can be rigid or flexible, hydrophilic or hydrophobic, symmetrical or unsymmetrical, have large surface area or small surface area, and interact or not with cell surface receptors. The molecular scaffold can be any of a variety of oligomers or polymers, including without limitation polyacrylamides, polyesters, polyethers, polymethacrylates, polyols, and polyamino acids and corresponding oligomers. Molecular scaffolds can in general be linear polymers, branched polymers or cross-linked polymers. Preferred molecular scaffolds are biocompatible. Molecular scaffolds prepared by ROMP methods, as illustrated in several formulas herein, are preferred. Molecular scaffolds can be hydrophobic or can be made to be more hydrophilic by substitution with polar substituents, such as -OH. The scaffold can be substituted, in general, with any groups that do not interfere with signal activity and which provide desirable chemical and physical properties.

25 The term "recognition element" or RE is used herein to refer to chemical or biochemical species, groups or structures that function for binding to cell receptors and in particular function or binding to cell surface receptors. RE are bonded to molecular scaffolds in the multivalent ligands of this invention. An RE can be a ligand for a cell receptor or a portion of such a ligand that is functional for receptor binding and that has been modified to allow its bonding to a molecular scaffold. An RE can be chemically identical to a cell receptor ligand or it may be modified from the ligand as a result of or to facilitate bonding to the scaffold.

5 The term "signal recognition element" or SRE is used herein to refer to chemical or biochemical species, groups or structure that function as chemical or biochemical signals (see below) and that are bonded into multivalent ligands of this invention. The SRE is typically a signal (group or molecule) that has been modified to allow its bonding into the multivalent ligand. An SRE can be chemically identical to a signal or it may be modified from the signal as a result of or to facilitate bonding to the scaffold. The SRE is preferably bonded into the multivalent ligand such that the signal function of the group is minimally affected. SREs are recognized by cells, typically by binding to a cell receptor, and thus are also REs. SREs, in contrast to REs, induce a response in or from the cell. The response may be an intracellular response, such as cell migration, and/or an intercellular response, such as the release of chemical species by the cell that function as chemical signals for other cells. Signal recognition is mediated by the presence of cell receptors on the cell surface to which the signal (or signal group) binds. Binding of signal (or SRE) alone may induce the biological response. Induction of the response may in some cases require presentation of multiple signals or (SRE). The biological response may in some cases be modulated by reorganization of receptors or clustering of receptors or the cell surface.

20 The term "chemical or biochemical signal" is used herein to refer to a particular chemical or biochemical species selected from various types (molecules, oligomers, moieties, groups etc.) that are recognized by a cell most typically by interaction with a cell surface receptor, and induce a biological response in the cell. A signal itself can induce the response on interaction with the cell or may only induce the response when multiple signals interact (e.g., when presented multivalently) with the cell. Signals can include the natural signals, which are those species found *in vivo* in a biological system to induce a response in or by a cell. Natural signals include, for example, naturally-occurring drugs, hormones, antigens, grow factors, cytokines, proteins, peptides, derivatized peptides (e.g., sulfated, phosphorylated, acylated, or N-formylated peptides), saccharides, derivatized saccharides (e.g., sulfated, acetylated, sialated), nucleic acids, various cell nutrients, epitopes and various small organic compounds (all of which may not represent mutually exclusive groups). Signals can also include chemical species that are found to mimic the function of natural chemical signals. These signal mimics are typically synthetic and can include, for example, synthetic drugs and various derivatives of naturally-occurring signals (e.g., peptoids and nucleic acid analogs or derivatives). Different cells can, of course, recognize different signals. Different cells may respond to the same or similar signals, with the same or with

different biological responses. A single cell may respond to a plurality of different signals to give the same or different biological response. Signals include, for example, chemoattractants and epitopes (antigenic determinants) which are not mutually exclusive groups. SREs bound to multivalent ligands can comprise a chemical or biochemical signal adapted for bonding to a molecular scaffold. SREs can include, among others, chemical and biochemical species that are chemoattractants, epitopes, cytokines, hormones and related substances.

*Smiley*  
*10/16*  
0985296-07001  
2007-10-16 09:52:00  
A chemoattractant is a chemical or biological signal toward which a cell migrates. The cell senses increasing concentrations of the chemoattractant and moves toward higher concentrations. Cell sensing mechanisms for chemoattractants are often very sensitive. Alternatively, cells may, in response to other signals, move to lower concentrations of signal. Bacterial cells migrate toward certain nutrients, such as glucose or galactose or amino acids, such as serine. Leukocytes (white blood cells) migrate toward, N-formyl peptides and other derivatized peptides, the activated component of CS (CSa), platelet-activating factor (PAF), leukotriene B4 (LTB4), or chemotactic cytokines (i.e., chemokines, including  $\alpha$ - and  $\beta$ -chemokines) (65). N-formylated peptides are products of bacterial protein synthesis and signal bacterial infection. The receptors for N-formylated peptides may also bind to other derivatized peptides such as N-acyl-peptides. Thus any ligand (which may include species that act as agonist or antagonists of receptor function) of a N-formylated peptide receptor may be employed for applications related to that receptor. Neutrophils, one type of leukocyte, are guided to the site of bacterial infection by sensing low levels of N-formylated peptides. Once at the site of infection phagocytosis can occur. A chemoattractant may induce biological responses in addition to migration or chemotaxis. For example, in various types of leukocytes, chemoattractants can induce the release of toxic species or the release of inflammatory cytokines, transcription factors and other chemical species which, in turn, function as chemical signals for other cells.

The term epitope is used generally herein to refer to any chemical species that functions as an antigenic determinant and most term generally includes all antigens. Epitopes are those parts of an antigen that combine with an antigen-binding site on an antibody molecule or on a lymphocyte (e.g., B cells and T cells) receptor. Binding of the epitope can, for example, stimulate antibody production or T cell responses. Epitopes may exhibit different levels of immunogenicity. Those that are more immunogenic than others and which dominant the overall antigenic response are designated immunodominant epitopes. Most

non-self proteins and many carbohydrates are antigens, so epitopes include, without limitation, proteins fragments (e.g., peptides) and carbohydrate fragments (e.g., saccharides and oligosaccharides). As used herein the term "self" as applied to antigen, epitope or cell is an entity that is recognized by an immune cell, a combination of immune cells or an immune system as self. The term "self" may also be applied other biological particles that are recognized as self by an immune cell, or cells or an immune system. Some anitgens, epitopes, cells and particles that are recognized as self are acutally foreign to the immune cell, cells or immune system, but are not so recognized. As used herein the term "foreign" as applied to antigen, epitope or cell is an entity that is recognized by an immune cell, a combination of immune cells or an immune system as foreign. Foreign is also any thing that is not recognized as self, i.e., non-self antigens, etc. The term "foreign" may also be applied to other biological particles that are recognized as foreign by an immune cell, or cells or an immune system. Some anitgens, epitopes, cells and particles that are recognized as foreign are acutally self to the immune cell, cells or immune system, but are not so recognized.

The term hapten takes its generally accepted meaning in the are as a small molecule, having at least one of the determinant groups of an antigen, that can combine with an antibody but is not immunogenic unless it acts in conjunction with a carrier molecule. Haptens include, among others, hemocyanins and nitro-sbustituted aromatic compounds, such as dinitrophenyl groups, trinitrobenzene sulphonyl groups, dinitrofluorophneyl groups.

The term anitbody as used herein is intended to encompass any protein or protein fragment that functions as an antibody and is specifically intended to include antibody fragmetns in cluding among others Fab fragments.

Lectin any of a large group of hemagglutinating proteins found principally in plant seeds. Certain lectins cause agglutination of erythrocytes of certain blood groups; others stimulate the proliferation of lymphocytes.

The term "biological system" is used generally herein to refer to any *in vivo* or *in vitro* system containing signal transduction elements, e.g., signal receptors and biochemical/biological elements for generating a response. A biological system typically contains at least one cell within any environment with which it interacts. A biological system in the context of the uses of multivalent ligands of this invention must contain at least one receptor which can interact with the ligand. In most applications of multivalent ligands, the biological system must contain at least one cell which can respond to the ligand. The response of a cell to the ligand occurs within the biological systems and as noted above may



1721  
cont

be an intracellular response, an intercellular response or both. The biological system can, for example, be a cell in a tissue, a cell in an organ or organism, a cell in a mixture of cells, a cell in a tissue culture, a cell in a tissue or biological fluid sample, and can include biological systems *in vivo* and *in vitro*.

5  
Sully #221

10

"Functional elements (FE)" are chemical or biochemical species (molecules, groups, moieties, etc.) that exhibit some biological or chemical function different from an RE or SRE. FE can, for example, provide reactive groups or latent reactive groups for attaching another chemical or biological group to a multivalent ligand. For example, an FE can be used to attach a multivalent ligand to a solid surface which may be useful for ligand purification or in applications to analytical or diagnostic assays. FE can be various detectable labels or reporter groups including fluorescent labels, radiolabels and high density labels such as gold particel bound to ligands (e.g., streptavidin labeled with gold particles). Multivalent ligands incorporating detectable labels or reporter groups can be used, for example, in various analytical or diagnostics assays. Of particular interest are multivalent ligands of this invention that are useful in visualization assays, e.g., for the detection of biological particles or molecules in microscopy applications. FE can also exhibit various biological functions, e.g., enzymatic function, ligand-binding function, etc., which may facilitate or enhance a selected application of a multivalent ligand.

20

Attachment of RE, SRE and/or FE can be facilitated by use of linker groups intervening between the molecular scaffold of the multivalent ligand and the signal group. Linker groups can be linear or branched, rigid or flexible, hydrophilic or hydrophobic as desired. One of ordinary skill in the art can select linkers from a variety of chemical species suitable for a given application. Further, one of ordinary skill in the art in view of methods and materials that are well known in the art can readily prepare multivalent ligands with

25

linkers having desirable properties.

30

Multivalent ligands of this invention can be used to modulate signal transduction in prokaryotic and eukaryotic organisms. The methods function in a variety of signal transduction processes. Prokaryote have a highly conserved intracellular signal transduction system, the two component system. The major components of this system are varying numbers of alternating histidine-aspartic acid kinase-mediated phosphorylation events, such as virulence, antibiotic resistance, response to environmental stress and sensing. The components of the two component system are highly conserved in prokaryotes. In contrast, eukaryotes appear to have very few two component systems for signal transduction. This

103  
5 orthogonality makes the two component signaling pathway a prime target for exploitation in therapeutic design for the control of bacterial infection. Major signal transduction systems in eukaryotes are mediated by G-protein-linked receptors and enzyme-linked receptors (including receptor guanylyl cyclases, receptor tyrosine kinases, tyrosine-kinase-associated receptors, receptor tyrosine phosphatases, and receptor serine/threonine kinases). The ability to modulate or regulate signal transduction in these pathways allows control over a wide variety of biological processes in eukaryotic cells and eukaryotic organism (including mammals and specifically humans) and provides significant opportunity for the design of therapeutics.

10  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995  
1000  
1005  
1010  
1015  
1020  
1025  
1030  
1035  
1040  
1045  
1050  
1055  
1060  
1065  
1070  
1075  
1080  
1085  
1090  
1095  
1100  
1105  
1110  
1115  
1120  
1125  
1130  
1135  
1140  
1145  
1150  
1155  
1160  
1165  
1170  
1175  
1180  
1185  
1190  
1195  
1200  
1205  
1210  
1215  
1220  
1225  
1230  
1235  
1240  
1245  
1250  
1255  
1260  
1265  
1270  
1275  
1280  
1285  
1290  
1295  
1300  
1305  
1310  
1315  
1320  
1325  
1330  
1335  
1340  
1345  
1350  
1355  
1360  
1365  
1370  
1375  
1380  
1385  
1390  
1395  
1400  
1405  
1410  
1415  
1420  
1425  
1430  
1435  
1440  
1445  
1450  
1455  
1460  
1465  
1470  
1475  
1480  
1485  
1490  
1495  
1500  
1505  
1510  
1515  
1520  
1525  
1530  
1535  
1540  
1545  
1550  
1555  
1560  
1565  
1570  
1575  
1580  
1585  
1590  
1595  
1600  
1605  
1610  
1615  
1620  
1625  
1630  
1635  
1640  
1645  
1650  
1655  
1660  
1665  
1670  
1675  
1680  
1685  
1690  
1695  
1700  
1705  
1710  
1715  
1720  
1725  
1730  
1735  
1740  
1745  
1750  
1755  
1760  
1765  
1770  
1775  
1780  
1785  
1790  
1795  
1800  
1805  
1810  
1815  
1820  
1825  
1830  
1835  
1840  
1845  
1850  
1855  
1860  
1865  
1870  
1875  
1880  
1885  
1890  
1895  
1900  
1905  
1910  
1915  
1920  
1925  
1930  
1935  
1940  
1945  
1950  
1955  
1960  
1965  
1970  
1975  
1980  
1985  
1990  
1995  
2000  
2005  
2010  
2015  
2020  
2025  
2030  
2035  
2040  
2045  
2050  
2055  
2060  
2065  
2070  
2075  
2080  
2085  
2090  
2095  
2100  
2105  
2110  
2115  
2120  
2125  
2130  
2135  
2140  
2145  
2150  
2155  
2160  
2165  
2170  
2175  
2180  
2185  
2190  
2195  
2200  
2205  
2210  
2215  
2220  
2225  
2230  
2235  
2240  
2245  
2250  
2255  
2260  
2265  
2270  
2275  
2280  
2285  
2290  
2295  
2300  
2305  
2310  
2315  
2320  
2325  
2330  
2335  
2340  
2345  
2350  
2355  
2360  
2365  
2370  
2375  
2380  
2385  
2390  
2395  
2400  
2405  
2410  
2415  
2420  
2425  
2430  
2435  
2440  
2445  
2450  
2455  
2460  
2465  
2470  
2475  
2480  
2485  
2490  
2495  
2500  
2505  
2510  
2515  
2520  
2525  
2530  
2535  
2540  
2545  
2550  
2555  
2560  
2565  
2570  
2575  
2580  
2585  
2590  
2595  
2600  
2605  
2610  
2615  
2620  
2625  
2630  
2635  
2640  
2645  
2650  
2655  
2660  
2665  
2670  
2675  
2680  
2685  
2690  
2695  
2700  
2705  
2710  
2715  
2720  
2725  
2730  
2735  
2740  
2745  
2750  
2755  
2760  
2765  
2770  
2775  
2780  
2785  
2790  
2795  
2800  
2805  
2810  
2815  
2820  
2825  
2830  
2835  
2840  
2845  
2850  
2855  
2860  
2865  
2870  
2875  
2880  
2885  
2890  
2895  
2900  
2905  
2910  
2915  
2920  
2925  
2930  
2935  
2940  
2945  
2950  
2955  
2960  
2965  
2970  
2975  
2980  
2985  
2990  
2995  
3000  
3005  
3010  
3015  
3020  
3025  
3030  
3035  
3040  
3045  
3050  
3055  
3060  
3065  
3070  
3075  
3080  
3085  
3090  
3095  
3100  
3105  
3110  
3115  
3120  
3125  
3130  
3135  
3140  
3145  
3150  
3155  
3160  
3165  
3170  
3175  
3180  
3185  
3190  
3195  
3200  
3205  
3210  
3215  
3220  
3225  
3230  
3235  
3240  
3245  
3250  
3255  
3260  
3265  
3270  
3275  
3280  
3285  
3290  
3295  
3300  
3305  
3310  
3315  
3320  
3325  
3330  
3335  
3340  
3345  
3350  
3355  
3360  
3365  
3370  
3375  
3380  
3385  
3390  
3395  
3400  
3405  
3410  
3415  
3420  
3425  
3430  
3435  
3440  
3445  
3450  
3455  
3460  
3465  
3470  
3475  
3480  
3485  
3490  
3495  
3500  
3505  
3510  
3515  
3520  
3525  
3530  
3535  
3540  
3545  
3550  
3555  
3560  
3565  
3570  
3575  
3580  
3585  
3590  
3595  
3600  
3605  
3610  
3615  
3620  
3625  
3630  
3635  
3640  
3645  
3650  
3655  
3660  
3665  
3670  
3675  
3680  
3685  
3690  
3695  
3700  
3705  
3710  
3715  
3720  
3725  
3730  
3735  
3740  
3745  
3750  
3755  
3760  
3765  
3770  
3775  
3780  
3785  
3790  
3795  
3800  
3805  
3810  
3815  
3820  
3825  
3830  
3835  
3840  
3845  
3850  
3855  
3860  
3865  
3870  
3875  
3880  
3885  
3890  
3895  
3900  
3905  
3910  
3915  
3920  
3925  
3930  
3935  
3940  
3945  
3950  
3955  
3960  
3965  
3970  
3975  
3980  
3985  
3990  
3995  
4000  
4005  
4010  
4015  
4020  
4025  
4030  
4035  
4040  
4045  
4050  
4055  
4060  
4065  
4070  
4075  
4080  
4085  
4090  
4095  
4100  
4105  
4110  
4115  
4120  
4125  
4130  
4135  
4140  
4145  
4150  
4155  
4160  
4165  
4170  
4175  
4180  
4185  
4190  
4195  
4200  
4205  
4210  
4215  
4220  
4225  
4230  
4235  
4240  
4245  
4250  
4255  
4260  
4265  
4270  
4275  
4280  
4285  
4290  
4295  
4300  
4305  
4310  
4315  
4320  
4325  
4330  
4335  
4340  
4345  
4350  
4355  
4360  
4365  
4370  
4375  
4380  
4385  
4390  
4395  
4400  
4405  
4410  
4415  
4420  
4425  
4430  
4435  
4440  
4445  
4450  
4455  
4460  
4465  
4470  
4475  
4480  
4485  
4490  
4495  
4500  
4505  
4510  
4515  
4520  
4525  
4530  
4535  
4540  
4545  
4550  
4555  
4560  
4565  
4570  
4575  
4580  
4585  
4590  
4595  
4600  
4605  
4610  
4615  
4620  
4625  
4630  
4635  
4640  
4645  
4650  
4655  
4660  
4665  
4670  
4675  
4680  
4685  
4690  
4695  
4700  
4705  
4710  
4715  
4720  
4725  
4730  
4735  
4740  
4745  
4750  
4755  
4760  
4765  
4770  
4775  
4780  
4785  
4790  
4795  
4800  
4805  
4810  
4815  
4820  
4825  
4830  
4835  
4840  
4845  
4850  
4855  
4860  
4865  
4870  
4875  
4880  
4885  
4890  
4895  
4900  
4905  
4910  
4915  
4920  
4925  
4930  
4935  
4940  
4945  
4950  
4955  
4960  
4965  
4970  
4975  
4980  
4985  
4990  
4995  
5000  
5005  
5010  
5015  
5020  
5025  
5030  
5035  
5040  
5045  
5050  
5055  
5060  
5065  
5070  
5075  
5080  
5085  
5090  
5095  
5100  
5105  
5110  
5115  
5120  
5125  
5130  
5135  
5140  
5145  
5150  
5155  
5160  
5165  
5170  
5175  
5180  
5185  
5190  
5195  
5200  
5205  
5210  
5215  
5220  
5225  
5230  
5235  
5240  
5245  
5250  
5255  
5260  
5265  
5270  
5275  
5280  
5285  
5290  
5295  
5300  
5305  
5310  
5315  
5320  
5325  
5330  
5335  
5340  
5345  
5350  
5355  
5360  
5365  
5370  
5375  
5380  
5385  
5390  
5395  
5400  
5405  
5410  
5415  
5420  
5425  
5430  
5435  
5440  
5445  
5450  
5455  
5460  
5465  
5470  
5475  
5480  
5485  
5490  
5495  
5500  
5505  
5510  
5515  
5520  
5525  
5530  
5535  
5540  
5545  
5550  
5555  
5560  
5565  
5570  
5575  
5580  
5585  
5590  
5595  
5600  
5605  
5610  
5615  
5620  
5625  
5630  
5635  
5640  
5645  
5650  
5655  
5660  
5665  
5670  
5675  
5680  
5685  
5690  
5695  
5700  
5705  
5710  
5715  
5720  
5725  
5730  
5735  
5740  
5745  
5750  
5755  
5760  
5765  
5770  
5775  
5780  
5785  
5790  
5795  
5800  
5805  
5810  
5815  
5820  
5825  
5830  
5835  
5840  
5845  
5850  
5855  
5860  
5865  
5870  
5875  
5880  
5885  
5890  
5895  
5900  
5905  
5910  
5915  
5920  
5925  
5930  
5935  
5940  
5945  
5950  
5955  
5960  
5965  
5970  
5975  
5980  
5985  
5990  
5995  
6000  
6005  
6010  
6015  
6020  
6025  
6030  
6035  
6040  
6045  
6050  
6055  
6060  
6065  
6070  
6075  
6080  
6085  
6090  
6095  
6100  
6105  
6110  
6115  
6120  
6125  
6130  
6135  
6140  
6145  
6150  
6155  
6160  
6165  
6170  
6175  
6180  
6185  
6190  
6195  
6200  
6205  
6210  
6215  
6220  
6225  
6230  
6235  
6240  
6245  
6250  
6255  
6260  
6265  
6270  
6275  
6280  
6285  
6290  
6295  
6300  
6305  
6310  
6315  
6320  
6325  
6330  
6335  
6340  
6345  
6350  
6355  
6360  
6365  
6370  
6375  
6380  
6385  
6390  
6395  
6400  
6405  
6410  
6415  
6420  
6425  
6430  
6435  
6440  
6445  
6450  
6455  
6460  
6465  
6470  
6475  
6480  
6485  
6490  
6495  
6500  
6505  
6510  
6515  
6520  
6525  
6530  
6535  
6540  
6545  
6550  
6555  
6560  
6565  
6570  
6575  
6580  
6585  
6590  
6595  
6600  
6605  
6610  
6615  
6620  
6625  
6630  
6635  
6640  
6645  
6650  
6655  
6660  
6665  
6670  
6675  
6680  
6685  
6690  
6695  
6700  
6705  
6710  
6715  
6720  
6725  
6730  
6735  
6740  
6745  
6750  
6755  
6760  
6765  
6770  
6775  
6780  
6785  
6790  
6795  
6800  
6805  
6810  
6815  
6820  
6825  
6830  
6835  
6840  
6845  
6850  
6855  
6860  
6865  
6870  
6875  
6880  
6885  
6890  
6895  
6900  
6905  
6910  
6915  
6920  
6925  
6930  
6935  
6940  
6945  
6950  
6955  
6960  
6965  
6970  
6975  
6980  
6985  
6990  
6995  
7000  
7005  
7010  
7015  
7020  
7025  
7030  
7035  
7040  
7045  
7050  
7055  
7060  
7065  
7070  
7075  
7080  
7085  
7090  
7095  
7100  
7105  
7110  
7115  
7120  
7125  
7130  
7135  
7140  
7145  
7150  
7155  
7160  
7165  
7170  
7175  
7180  
7185  
7190  
7195  
7200  
7205  
7210  
7215  
7220  
7225  
7230  
7235  
7240  
7245  
7250  
7255  
7260  
7265  
7270  
7275  
7280  
7285  
7290  
7295  
7300  
7305  
7310  
7315  
7320  
7325  
7330  
7335  
7340  
7345  
7350  
7355  
7360  
7365  
7370  
7375  
7380  
7385  
7390  
7395  
7400  
7405  
7410  
7415  
7420  
7425  
7430  
7435  
7440  
7445  
7450  
7455  
7460  
7465  
7470  
7475  
7480  
7485  
7490  
7495  
7500  
7505  
7510  
7515  
7520  
7525  
7530  
7535  
7540  
7545  
7550  
7555  
7560  
7565  
7570  
7575  
7580  
7585  
7590  
7595  
7600  
7605  
7610  
7615  
7620  
7625  
7630  
7635  
7640  
7645  
7650  
7655  
7660  
7665  
7670  
7675  
7680  
7685  
7690  
7695  
7700  
7705  
7710  
7715  
7720  
7725  
7730  
7735  
7740  
7745  
7750  
7755  
7760  
7765  
7770  
7775  
7780  
7785  
7790  
7795  
7800  
7805  
7810  
7815  
7820  
7825  
7830  
7835  
7840  
7845  
7850  
7855  
7860  
7865  
7870  
7875  
7880  
7885  
7890  
7895  
7900  
7905  
7910  
7915  
7920  
7925  
7930  
7935  
7940  
7945  
7950  
7955  
7960  
7965  
7970  
7975  
7980  
7985  
7990  
7995  
8000  
8005  
8010  
8015  
8020  
8025  
8030  
8035  
8040  
8045  
8050  
8055  
8060  
8065  
8070  
8075  
8080  
8085  
8090  
8095  
8100  
8105  
8110  
8115  
8120  
8125  
8130  
8135  
8140  
8145  
8150  
8155  
8160  
8165  
8170  
8175  
8180  
8185  
8190  
8195  
8200  
8205  
8210  
8215  
8220  
8225  
8230  
8235  
8240  
8245  
8250  
8255  
8260  
8265  
8270  
8275  
8280  
8285  
8290  
8295  
8300  
8305  
8310  
8315  
8320  
8325  
8330  
8335  
8340  
8345  
8350  
8355  
8360  
8365  
8370  
8375  
8380  
8385  
8390  
8395  
8400  
8405  
8410  
8415  
8420  
8425  
8430  
8435  
8440  
8445  
8450  
8455  
8460  
8465  
8470  
8475  
8480  
8485  
8490  
8495  
8500  
8505  
8510  
8515  
8520  
8525  
8530  
8535  
8540  
8545  
8550  
8555  
8560  
8565  
8570  
8575  
8580  
8585  
8590  
8595  
8600  
8605  
8610  
8615  
8620  
8625  
8630  
8635  
8640  
8645  
8650  
8655  
8660  
8665  
8670  
8675  
8680  
8685  
8690  
8695  
8700  
8705  
8710  
8715  
8720  
8725  
8730  
8735  
8740  
8745  
8750  
8755  
8760  
8765  
8770  
8775  
8780  
8785  
8790  
8795  
8800  
8805  
8810  
8815  
8820  
8825  
8830  
8835  
8840  
8845  
8850  
8855  
8860  
8865  
8870  
8875  
8880  
8885  
8890  
8895  
8900  
8905  
8910  
8915  
8920  
8925  
8930  
8935  
8940  
8945  
8950  
8955  
8960  
8965  
8970  
8975  
8980  
8985  
8990  
8995  
9000  
9005  
9010  
9015  
9020  
9025  
9030  
9035  
9040  
9045  
9050  
9055  
9060  
9065  
9070  
9075  
9080  
9085  
9090  
9095  
9100  
9105  
9110  
9115  
9120  
9125  
9130  
9135  
9140  
9145  
9150  
9155  
9160  
9165  
9170  
9175  
9180  
9185  
9190  
9195  
9200  
9205  
9210  
9215  
9220  
9225  
9230  
9235  
9240  
9245  
9250  
9255  
9260  
9265  
9270  
9275  
9280  
9285  
9290  
9295  
9300  
9305  
9310  
9315  
9320  
9325  
9330  
9335  
9340  
9345  
9350  
9355  
9360  
9365  
9370  
9375  
9380  
9385  
9390  
9395  
9400  
9405  
9410  
9415  
9420  
9425  
9430  
9435  
9440  
9445  
9450  
9455  
9460  
9465  
9470  
9475  
9480  
9485  
9490  
9495  
9500  
9505  
9510  
9515  
9520  
9525  
9530  
9535  
9540  
9545  
9550  
9555  
9560  
9565  
9570  
9575  
9580  
9585  
9590  
9595  
9600  
9605  
9610  
9615  
9620  
9625  
9630  
9635  
9640  
9645  
9650  
9655  
9660  
9665  
9670  
9675  
9680  
9685  
9690  
9695  
9700  
9705  
9710  
9715  
9720  
9725  
9730  
9735  
9740  
9745  
9750  
9755  
9760  
9765  
9770  
9775  
9780  
9785  
9790  
9795  
9800  
9805  
9810  
9815  
9820  
9825  
9830  
9835  
9840  
9845  
9850  
9855  
9860  
9865  
9870  
9875  
9880  
9885  
9890  
9895  
9900  
9905  
9910  
9915  
9920  
9925  
9930  
9935  
9940  
9945  
9950  
9955  
9960  
9965  
9970  
9975  
9980  
9985  
9990  
9995  
10000  
10005  
10010  
10015  
10020  
10025  
10030  
10035  
10040  
10045  
10050  
10055  
10060  
10065  
10070  
10075  
10080  
10085  
10090  
10095  
10100  
10105  
10110  
10115  
10120  
10125  
10130  
10135  
10140  
10145  
10150  
10155  
10160  
10165  
10170  
10175  
10180  
10185  
10190  
10195  
10200  
10205  
10210  
10215  
10220  
10225  
10230  
10235  
10240  
10245  
10250  
10255  
10260  
10265  
10270  
10275  
10280  
10285  
10

cell. If FE has a biological function, the multivalent ligand targets the cell with that function. Multivalent ligands that contain a plurality of RE (SRE or both) can function in macromolecular assembly which need not involve any biological signaling function. In such applications, the multivalent ligand need not contain any SRE, the multivalent ligand need only contain more than one recognition element for binding to a cell surface receptor (a recognition element, RE) and preferably a plurality of REs. In such applications, the multivalent ligands directly or indirectly bind to more than one cell resulting in cell aggregation. Cell aggregation may itself trigger a biological response (e.g., the release of signal molecules by a cell), but need not. Multivalent ligands can indirectly cause cell aggregation by binding to a plurality of biochemical species, such as lectins (e.g., Concanavalin A) which in turn bind to cells resulting in cell aggregation. The effect of a multivalent ligand on indirect cell aggregation will be dependent upon the valency of the ligand and on the relative concentrations of the multivalent ligand to the species that causes cell aggregation. At higher concentrations of multivalent ligands with higher valency, binding sites on the species that causes cell aggregation may be saturated inhibiting cell aggregation. At lower concentrations of multivalent ligand, free binding sites will remain and cell aggregation can occur and can be enhanced by the multivalent ligand. Thus, multivalent ligands of this invention can be selectively designed to inhibit or to facilitate cell aggregation.


Multivalent ligands functioning for macromolecular assembly can be useful in a variety of applications, particularly those based on cell aggregation, including, but not limited to diagnostic assays, cancer therapy, and pathogen clearance.

The reorganization of receptors on cell surfaces is involved in many important biological reactions including cell migration, adhesion, and the formation of cell to cell junctions. Multivalent ligands of this invention and in particular those ligands which can span the distance between receptors, as discussed above, can be used to reorganize receptors and to modulate response due to the individual signal interactions with the receptors. Reorganization of receptors on the cell surface includes without limitation: changing the relative positions of different cell receptors on the surface, lateral movement of receptors on the surface, the localization of receptors to different sites on the cell surface, changes in the proximity of signal transduction machinery associated with receptors, changes in the proximity of features of the intracellular matrix associated with receptors, changes in the

proximity of receptors, clustering of receptors, changes in conformation of receptors, and initiation of receptor-receptor interactions.

10 In specific embodiments, linear multivalent ligands of this invention are prepared by ring opening metathesis polymerization (ROMP), see for example (54). This method has been used to prepare multivalent inhibitors of cell functions (27, 28). The ROMP methods have been described in more detail in U.S. patent 5,587,442 relating to multivalent ligands that are polyglycomers. Improvement of ROMP methods for generating block polymers (and oligomers) and for introducing end-groups on ROMP polymers ( and oligomers) have been described in U.S. patent applications 09/335,420 and 09/336,121, both filed June 17, 1999. (These U.S. patent documents are incorporated by reference herein in their entirety particularly for the description of ROMP methods). Scheme 6 illustrates exemplary methods for modification of ROMP backbones, which can be applied in combination with synthetic methods described in the above listed patents and patent applications to synthesize multivalent ligands of this invention. Scheme 6 illustrates a diimide reduction (23, 98, 99) which can be employed to reduce double bonds in ROMP scaffold backbones. Scheme 6 also illustrates the substitution of ROMP scaffold backbones with OH groups using OsO<sub>4</sub> catalyzed dihydroxylation (100, 101). Those of ordinary skill in the art can prepare multivalent ligands of this invention, particularly those specifically exemplified in formulas herein, employing the descriptions herein and methods that are well known in the art.

Multivalent ligands of this invention prepared by ROMP are exemplified by the general structure:



wherein:

n is an integer that is 2 or more and represents the number of repeating unit in parentheses that are in the ligand;

the dashed lines indicate optional double bonds;

“BB” represents the backbone repeating unit, which may be cyclic or acyclic, and may be the same or different in a random or block arrangement where the wavy lines indicate that the BB repeating unit can be in a cis or trans configuration in the backbone;

R<sup>1</sup> and R<sup>2</sup>, can be H, an organic group, an FE group or the groups: -L-RE- or -L-SRE- wherein FE is a functional element other than an RE or an SRE, L represents an optional linker group, RE is a recognition element and SRE is a signal recognition element;

R<sup>4</sup> and R<sup>5</sup> are H, or an organic group;


R<sup>6</sup> and R<sup>7</sup> are H, an organic group or an end-group;

Z, independently of other Z in the polymer, is H, OH, OR<sup>8</sup>, SH, a halide (F, Br, Cl, I), NH<sub>2</sub> or N(R<sup>8</sup>)<sub>2</sub> where R<sup>8</sup> is H or an organic group or Z is absent when there is a double bond at the carbon to which A is attached.

The integer n is the average number of repeating units in the polymer. Typically n can range up to about 10,000, but there is no practical limit. Preferably the number of repeating units in the multivalent ligands of this invention is defined and can range generally from 2 up to several hundred or several thousand. Preferred multivalent ligands will have n that ranges from 10 to about 100. Multivalent ligands of this invention also include those in which n ranges from 10 to about 25, in which n is 25 or more and those in which n is 50 or more. ROMP can provide polymers of varying average lengths (i.e., varying degree of polymerization, DP) depending on the monomer to ROMP catalyst (i.e., initiator) ratios. The length of all polymers referred to herein (i.e., n or n+m, below) is the length predicted by the monomer to initiator ratios used in the polymerization reaction.

BB can be alkyl, cycloalkyl, cycloalkenyl, and one or more CH<sub>2</sub> groups in the BB moiety can be replaced with -O-, -S-, -NR<sup>9</sup>-, or -CO-, where R<sup>9</sup> is H or an organic group. Preferred BB have 10 or fewer carbon atoms.

Exemplary BB repeating units include among others:



where Y can be -O-, -S-, -NR<sup>8</sup>, or -CH<sub>2</sub>-  
and there is an optional double bond  
indicated by the dashed line

(CH<sub>2</sub>)<sub>q</sub>

where q is 1 to about 10

(CH<sub>2</sub>)<sub>p</sub>-NH-CO-(CH<sub>2</sub>)<sub>m</sub>

where p is 0 to about 10 and m is 0 to about 10

RE is a recognition element as discussed above that can be any of a variety of chemical or biochemical species that are recognized by and which selective bind to cell receptors, particularly, transmembrane receptors and cell surface receptors. SRE is a signal recognition element as discussed above that can be any of a variety of chemical or biochemical species that are recognized by one or more cells and which induce a biological response by the cell; "L" is an optional linker group that can provide functional groups for covalent bonding of the RE, SRE or FE to the polymer (oligomer) backbone. FE is a chemical or biochemical functional group other than an SRE, as discussed above. Other examples of ROM scaffolds are illustrated in Schemes 2 and 3.

The multivalent ligand of the above formula contains up to n RE, SRE or both. In specific embodiments all of the monomers carry an RE or SRE (the number of RE + SRE is n). In other specific embodiments, regions of spacer monomers that do not carry RE or SRE intervene between regions of monomers that carry RE or SRE. The RE and SRE attached to different monomers may be the same or different. In one embodiment, RE or SRE throughout the multivalent ligand are all the same. In another embodiment, the multivalent ligand contains more than one type of RE or SRE. In a specific embodiment, the multivalent ligand contains two different types of RE or SRE or an RE and an SRE. In this embodiment, the RE and SRE are non-randomly positioned in the ligand. Preferably monomers carrying the same RE or SRE are grouped into blocks (as in block polymers) within the multivalent ligand and spacer monomers are optionally positioned between blocks. In other embodiments, R<sup>1</sup> and R<sup>2</sup> together can form an RE or SRE.

RE and SRE are attached to the polymer (oligomer) backbone such that they substantially retain their function for binding cell receptors or as signals, respectively. For a given RE or SRE there may be several ways in which it can be bonded into the multivalent

ABO

5 ligand, each of which may result in RE that are different in binding affinity or SRE that are different either in binding affinity or in the level or type of response induced. For example, a peptide signal may be bonding through its N-terminus, through its C-terminus or via an amino acid side group, such as through a lysine side group. The site of attachment of an RE or SRE to the multivalent ligand is preferably selected to minimize loss of binding function (RE) or to minimize loss of signal function (SRE) or alternatively site of attachment may be selected to maximize signal function (SRE). An RE or SRE may nevertheless exhibit properties that are different from free ligands or free signals (e.g., the binding affinity of an SRE for a cell receptor may be different from that of free signal from which it was derived or which it mimics), but which do not destroy the function of an RE as a ligand or an SRE as a signal. RE can include a variety of known cell receptor ligands and in particular can include lectins. SRE can specifically include monosaccharides (e.g., glucose, galactose), disaccharides, polysaccharides (greater than 2 sugar residues), derivatized saccharides (e.g., acylated, sialated), peptides, derivatized peptides (e.g., N-formyl peptides), peptoids, various chemoattractants, and various epitopes. Note that a particular chemical or biological species may function as an RE with one type or kind of cell and as an SRE with another type or kind of cell.

10

00015296  
"C" 1001

25 The linker can provide for spacing of the RE, SRE or FE group(s) from the backbone or can provide for structural flexibility. Linkers may be the same or different on different monomers in the polymer. Linkers that are used in a monomeric scaffold to bond to RE, SRE or FE can also be all the same or different. In a given multivalent ligand carrying one type of RE or SRE group, the linker is preferably the same throughout the polymer. Linkers are generally selected so that they are compatible with the intended application of the multivalent ligand and to avoid interference with the function of signal groups. The linker is preferably linear and preferably ranges in length from 1 to about 20 atoms. The linker may contain alicyclic groups (such as a cyclohexyl group). The linker can be an alkyl chain carrying functional groups for bonding to the backbone of the ligand and to the signal. The linker can also be an ether, ester, ketone, amine, amide or thioether chain. In a specific embodiment, the linker can be described as an linear alkyl chain having from 1 to about 20 carbon atoms in length in which one or more non-neighboring CH<sub>2</sub> groups are optionally replaced with an -O-, -S-, -NH-, -NR<sup>10</sup>-, -CO-, -NH-CO-, -O-CO-, -C=C-, or -C≡C- group, where R<sup>10</sup> is an alkyl or aryl group. Linker CH<sub>2</sub> groups can be substituted with halogens, alkoxy, or alkyl groups. In the absence of a linker group, the ROMP backbone or the signal group itself must provide

30

A31 the functionality for covalent bonding of the signal to the backbone. Exemplary linkers include those illustrated in Scheme 3.

Sul A32  
5 R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> can be organic groups. Organic groups include without limitation alkyl groups, alkenyl groups, and aryl groups as well as substituted alkyl, alkenyl and aryl groups. Substituents for alkyl, alkenyl and aryl groups include halogens (F, Cl, Br, I), -CN, -NO<sub>2</sub>, -OH, -SH, -NH<sub>2</sub>, -N(R<sup>10</sup>)<sub>2</sub>, -SR<sup>10</sup> and -OR<sup>10</sup> where R<sup>10</sup> is an alkyl or aryl group. Aryl groups may also contain alkyl or alkenyl substituents. Organic groups will typically have from 1 to about 20 carbon atoms, and preferably have 1 to about 10 carbon atoms. Alkyl groups may be straight-chain, branched or cyclic (or contain portions that are cyclic). One or more non-neighboring -CH<sub>2</sub>- groups in an alkyl or alkenyl group can be replaced with -O-, -S-, -NH- or -NR<sup>10</sup>, where R<sup>10</sup> is an alkyl or aryl group.

10 R<sup>6</sup> and R<sup>7</sup> can be end-groups, such as those described in U.S. patent application 09/336,121 filed June 17, 1999 which is incorporated in its entirety herein for description of methods of synthesis of multivalent ligands having end-groups using ROMP methods. End-groups can include a latent reactive group or a non-reactive functional group as described in the cited patent application. The presence of a latent reactive group would allow for later functionalization of a polymer multivalent ligand at an end-group. End-groups can contain functionality for binding to solid surfaces. The end-group may itself be a linkage to a solid support material. Latent reactive groups include: azides, a nitro group, a disulfide, a cyano group, an acetal group, a ketal, a carbamate, a thiocyanate, an activated ester, or an activated acid (activated esters and acids are those containing good leaving groups that are activated in particular for nucleophilic attack). Non-reactive end-groups include natural products or analogs thereof (e.g., biotin), metal chelators (e.g., nitrilotriacetic acid), metals (e.g., Zn), and fluorescent labels (amide derived BODIPYL FL EDA which is 4,4-difluoro-5,7-dimethyl-4-boro-3a,4a-diaza-s-indacene-3-propionyl ethylenediamine). End-groups can include FE.


25 The multivalent ligand optionally contains one or more functional elements that are not SRE. Preferred multivalent ligands contain significantly fewer FE compared to SRE. FE can be or contain any of the reactive or non-reactive groups listed above or described in U.S. patent application 09/336,121 filed June 17, 1999 as "end-groups". FE can also have enzymatic or other protein function.

30 Sul A33 When prepared by the ROMP methods, such as those described in U.S. patent applications 09/335,420 and 09/336,121, both filed June 17, 1999 (which are incorporated by reference herein in their entirety for methods of synthesis of multivalent ligands), R<sup>4</sup> and R<sup>5</sup>




are derived from the metal carbene catalysts, i.e., they are substituents on the metal carbene carbon of the metal carbene catalyst and in specific embodiments are H and a phenyl group. When using ROMP,  $R^6$  and  $R^7$  are typically derived from the capping agent, i.e., are the substituents on the electron rich alkene capping agent, such as hydrogen in the case of ethyl vinyl ether.

In a specific embodiment multivalent ligands of this invention include those of formula:




wherein BB,  $R^{1-2}$ , and  $R^{4-7}$  are as defined above. In specific embodiments, one of  $R^1$  or  $R^2$  is H and the other is L-RE. In specific embodiments, one or  $R^1$  or  $R^2$  is H and the other is L-SRE. In specific embodiments, RE is a lectin or a cell receptor ligand that is comprised within a lectin. In specific embodiments, SRE is a monosaccharide, a disaccharide or a relatively short saccharide having up to about 10 sugar residues. In other specific embodiments, SRE is a peptide or a derivatized peptide (e.g., an N-formyl peptide).

In another specific embodiment the invention relates to multivalent ligands of the formula:




wherein the dashed line indicates an optional double bond and wherein Y, independently of Y in other monomers,  $R^{1-2}$ , independent of  $R^{1-2}$  in other monomers, and  $R^{4-7}$  are as defined above. In specific embodiments, Y is  $-CH_2-$ . In specific embodiments, one of  $R^1$  or  $R^2$  is H and the other of  $R^1$  or  $R^2$  is  $-L-RE$ . In specific embodiments, one of  $R^1$  or  $R^2$  is H and the other of  $R^1$  or  $R^2$  is  $-L-SRE$ .  $R^1$  and  $R^2$  together may form an  $-L-RE$  or  $-L-SRE$ . In yet other specific embodiments, SRE is a peptide or derivatized peptide. When no double bond is present the ring carbons typically carry addition hydrogens, but may be substituted with other groups, such as alkyl groups having 1-6 carbon atoms or halides that do not interfere with the function of any  $R^1$  or  $R^2$  group.

10 In yet another specific embodiment the invention relates to multivalent ligands of the formulas:




or




in which m is the number of monomers carrying a first SRE (SRE<sup>1</sup>) and n is the number of monomers carrying a second SRE (SRE<sup>2</sup>). L<sup>1</sup> and L<sup>2</sup> are linkers as described above which may be the same or different. All other variables are as defined in earlier formulas and dashed lines indicating optional double bonds. Both m and n are integers that can range most generally from 1 up to about 10,000, but which more typically will range from 1 to several hundred or several thousand. The value of m may be the same as or different from that of n. In preferred ligands, n + m ranges from 5 or more up to about 200. Multivalent ligands of this invention include those in which n + m ranges between about 10 and 25, those in which n + m is 25 or more, those in which n + m is 50 or more, and those in which n + m is 100 or more.

Other exemplary multivalent ligands include those of the formulas:



wherein n, m and p are integers with a value greater than 3 and other variables are as defined above and



wherein n, m, p and x are integers each of which has a value greater than 1 and all other variables are as defined above. Multivalent ligands of these formulas can contain multiple blocks of

monomer regions having the same RE or SRE. Multivalent ligands of these formulas can contain multiple blocks of monomer regions one RE or SRE and multiple blocks of monomer regions containing another RE or SRE. Multivalent ligands of these formulas can also contain multiple blocks of monomer regions carrying RE or SRE with intervening spacer regions that carry no RE or SRE.

The multivalent ligands of this invention are useful in methods for controlling or modulating the effect of chemical signals in a biological system. Applications of multivalent ligands to bacterial and eukaryotic chemotaxis, to migration of leukocytes (particularly neutrophils), to immune responses of B-cells and T-cells, to cell aggregation, and to signaling of apoptosis are exemplified herein below.

Multivalent ligands of this invention which carry bacterial chemoattractants can be employed to disrupt colonization and biofilm formation by bacteria. Chemotaxis is a virulence factor which facilitates bacterial colonization of its host. Disruption of colonization of host tissue prevents host-bacterial interactions, prevents colonization and inhibits or retards infection. The methods of this invention can be applied specifically to disruption of colonization, for example, by *Staphylococcus aureus* (for treatment of staph infections) and *Vibrio cholerae* (for treatment of cholera). One bacterial survival mechanism involves the formation of microcommunities with functional heterogeneity (biofilms). Biofilm formation and maintenance are regulated by soluble small molecule-based factors. These factors control signal transduction pathways that allow bacteria to sense their environment and conversions to biofilm formation are mediated by two-component signaling systems. Disruption of biofilm formation renders bacteria more susceptible to host defenses or to antibiotic treatment and can inhibit or retard infection. Multivalent ligands which disrupt biofilm formation can be particularly useful in preventing or treating infections of the lung, for example for treating or preventing lung infection by the opportunistic pathogen *Pseudomonas aeruginosa*. Infection by this organism is a leading cause of death in patients with cystic fibrosis. Another mechanism for bacterial survival is induction of a virulence response upon increased bacterial cell density. This virulence response is induced by the release of soluble factors when increased cell density is sensed. Disruption of the responses of bacteria to increased cell density by multivalent ligands of this invention can be used to control bacterial virulence, for example, this method is applicable to the control of virulence of *Staphylococcus aureus*.

Multivalent ligands of this invention can in similar ways be employed to disrupt infection by eukaryotic pathogens and parasites, including among others, *Trypanosoma cruzi* (chuga disease) *Trypanosoma brucei* (sleeping sickness), tapeworms, hookworms, and *Plasmodium falciparum* (malaria).

5           The multivalent ligands of this invention can be used to modulate immune response toward epitopes and antigens (e.g. by modulating the immunogenicity of these species). For example, multivalent ligands can be designed to stimulate or inhibit leukocyte responses, including migration. Stimulation of such response can be used to enhance recognition of non-self cells for clearance and treat infection. Multivalent ligands can also be designed to  
10           modulate the activation and/or deactivation of B-cells or T-cells in response to chemical signals to improve and enhance desired immune response. B-cells and T-cells can be treated with multivalent ligands of this invention *in vitro*, *in vivo* and *ex vivo*.

          Autoimmune diseases involve aberrant function of a cell signal recognition process in which self cells are incorrectly marked for clearance. Multivalent ligands of this invention  
15           which modulate cell responses of immune system cells to epitopes can be employed to inhibit or attenuate autoimmune disorders. In a specific embodiment, ligands carrying self epitopes mistakenly recognized as "non-self" and certain B-cell or T-cell epitopes can be employed in a tolerization process to ameliorate autoimmune responses.

          The multivalent ligands of this invention also have application to the treatment of undesired cell proliferation (cancer) and undesired cell migration (metastasis). Cancer cells  
20           have distinct surface features (e.g., epitopes) that distinguish them from non-cancer cells. The multivalent ligands of this invention can be designed to promote recognition of cancer-specific epitopes as non-self cells by the immune system such that cancer cells are cleared by the immune system. Multivalent ligands carrying cancer cell epitopes and B-cell or T-cell  
25           epitopes can be employed in a sensitization process to promote clearance of the cancer cells. Cancer metastasis is deviant cell migration. The movement, adhesion, and junction formation of cancer cells are mediated, at least in part, by interaction of cancer cells with the multivalent extracellular matrix. Multivalent ligands can be designed to inhibit or prevent movement, adhesion and junction formation and thus inhibit metastasis.

30           This invention provides pharmaceutical and therapeutic compositions comprising multivalent ligands with SRE groups selected to provide therapeutic benefit in combination with a pharmaceutically acceptable carrier or excipient adapted for use in human or veterinary medicine. The multivalent ligands may be combined with each other to achieve a

desired pharmaceutical response or administered in combination with other known drugs or therapeutic agents, including without limitation antibacterial and other antimicrobial agents. The multivalent ligand is present in the pharmaceutical compositions in an amount, or in combination with other ligands in a combined amount, sufficient to obtain the desired therapeutic benefit. The carrier or excipient is selected as is known in the art for compatibility with the desired means of administration, for compatibility with the selected multivalent ligand(s) and to minimize detrimental effects to the patient.

This invention is also directed to pharmaceutically acceptable esters and salts of the multivalent ligands of various formulas and structures described herein. Acid addition salts are prepared by contacting compounds having appropriate basic groups therein with an acid whose anion is generally considered suitable for human or animal consumption.

Pharmacologically acceptable acid addition salts include but are not limited to the hydrochloride, hydrobromide, hydroiodide, sulfate, phosphate, acetate, propionate, lactate, maleate, malate, succinate, and tartrate salts. All of these salts can be prepared by conventional means by reacting, for example, the selected acid with the selected basic compound. Base addition salts are analogously prepared by contacting compounds having appropriate acidic groups therein with a base whose cation is generally considered to be suitable for human or animal consumption. Pharmacologically acceptable base addition salts, include but are not limited to ammonium, amine and amide salts.

Pharmaceutically acceptable esters of compounds of this invention are prepared by conventional methods, for example by reaction with selected acids. Pharmaceutically acceptable esters include but are not limited to carboxylic acid esters  $R^E\text{COO-D}$  (where D is a cationic form of a compound of this invention and where  $R^E$  is H, alkyl or aryl groups).

This invention is also directed to prodrugs of multivalent ligands and derivatives which on being metabolized will result in any of the ligands of this invention. Labile substituents may be protected employing conventional and pharmaceutically acceptable protecting groups removable on metabolism. Pharmaceutically active compounds may be derivatized by conventional methods to provide for extended metabolic half-life, to enhance solubility in a given carrier, to provide for or facilitate slow-release or timed-release or enhance or affect other drug delivery properties.

The multivalent ligands according to the invention may be formulated for oral, buccal, parenteral, topical or rectal administration. In particular, the ligands according to the invention may be formulated for injection or for infusion and may be presented in unit dose

form in ampules or in multidose containers with an added preservative. The compositions may take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g. sterile, pyrogen-free water, before use.

The pharmaceutical compositions according to the invention may also contain other active ingredients, such as antimicrobial agents, or preservatives. In general, pharmaceutical compositions of this invention can contain from 0.001-99% (by weight) of one or more of a multivalent ligands described herein.

For administration by injection or infusion, the daily dosage as employed for treatment of an adult human of approximately 70 kg body weight will range from 0.2 mg to 10 mg, preferably 0.5 to 5 mg, which can be administered in 1 to 4 doses, for example, depending on the route of administration and the clinical condition of the patient. These formulations also include formulations in dosage units. This means that the formulations are present in the form of a discrete pharmaceutical unit, for example, as tablets, dragees, capsules, caplets, pills, suppositories or ampules. The active compound content of each unit is a fraction or a multiple of an individual dose. The dosage units can contain, for example, 1, 2, 3 or 4 individual doses for  $\frac{1}{2}$ ,  $\frac{1}{3}$  or  $\frac{1}{4}$  of an individual dose. An individual dose preferably contains the amount of active compound which is given in one administration and which usually corresponds to a whole, one half, one third or one quarter of a daily dose. The magnitude of a prophylactic or therapeutic dose of a particular multivalent ligand will, of course, vary with the nature of the severity of the condition to be treated, the particular ligand compound and its route of administration. It will also vary according to the age, weight and response of the individual patient.

The compounds of the present invention are preferably formulated prior to administration. The present pharmaceutical formulations are prepared by known procedures using well-known and readily available ingredients. In making the compositions of the present invention, the active ingredient will usually be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material which acts as a vehicle, excipient or medium for the active ingredient. The compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium),

ointments containing for example up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions and sterile packaged powders.

Some examples of suitable carriers, excipients, and diluents include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, methyl cellulose, methyl and propylhydroxybenzoates, talc, magnesium stearate and mineral oil. The formulations can additionally include lubricating agents, wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavoring agents. The compositions of the invention may be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures well known in the art.

The compositions are preferably formulated in a unit dosage form, each dosage containing from about 0.5 to about 150 mg, more usually about 0.1 to about 10 mg, of the active ingredient. The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical carrier.

The invention is further directed to therapeutic methods that comprise the step of administering a pharmaceutical composition of this invention to an individual that can derive therapeutic benefit from the compositions.

Multivalent ligands of this invention can also be employed in non-therapeutic applications, for example, to prevent or inhibit biofouling in a selected environment or to remove undesired cells from a selected environment. Compositions of this invention for use in such non-therapeutic comprise one or more multivalent ligands of this invention in an amount or in a combined amount effective for obtaining a desired function, e.g., effective for affecting bacterial or microbial chemotaxis or effective for aggregating cells in a sample or a biological system. Compositions can be formulated using any appropriate solvent or carrier system which may be an aqueous solution, a lyophilized or a spray-dried material so long as desired function is maintained.

The following examples, further illustrate and further describe the invention, but are in no way intended to limit the invention.



## THE EXAMPLES

### Modulation of Bacterial Chemotaxis

09815296-073001  
TIME 20 " 96251860

The molecular events leading to bacterial chemotaxis have been well studied, and the process has served as a general model for receptor-mediated responses [29-32]. During chemotaxis in *Escherichia coli*, chemoattractants, such as sugars and amino acids, and chemorepellents are recognized by specific receptors at the bacterial plasma membrane [33]. For these investigations of multivalent ligand activity, galactose was selected as a model chemoattractant. The related compound,  $\beta$ -methyl galactopyranoside, is also a chemoattractant, indicating that the attachment of substituents at the anomeric position of galactose does not abolish its chemotactic activity [34]. This observation suggests that galactose residues can be tethered through an anomeric linker to create a multivalent display. For galactose-mediated signaling, the saccharide must bind to the soluble periplasmic glucose/galactose-binding protein (GGBP), which, in turn, interacts with the galactose-sensing chemoreceptor, Trg [34, 35]. Galactose-GGBP binding to Trg initiates a signaling pathway that results in reversal of the direction of flagellar spin, allowing the bacteria to swim towards the nutrient [29, 30, 36].

Bacterial chemotaxis requires an extremely sensitive sensing system with a broad dynamic range. Through their chemoreceptors, bacteria can detect very small changes in ligand concentration over many orders of magnitude [37, 38]. A recent mathematical model proposed by Bray *et al.* to explain this remarkable feature suggests signal transduction is regulated by changes in lateral clustering of the chemoreceptors [39-41]. In this model, clusters of bacterial chemoreceptors exchange ligand binding information, such that receptor clusters are more active in signal generation than individual receptors [39, 41]. Multivalent ligands of this invention having distinct valencies can differentially reorganize the receptors and thus control lateral receptor organization may result in modulation of the chemotactic response.

Galactose-bearing ligands 1-4 of varying defined valencies were generated using ROMP methods (Scheme 1). The galactose residues in the multivalent ligands are tethered to the molecular scaffold (polymer backbone) via a short linker. The interaction of monomer 1 was at least as favorable as that of galactose in an *in vitro* binding assay, thus the attachment of the linker did not prevent galactose binding to purified GGBP.

Ligands functionalized with galactose such as monovalent ligand 1 and multivalent ligand 3, also serve as attractants *in vivo*. This was demonstrated by monitoring the

behavioral response of *E. coli* to these ligands. The locomotion behavior of *E. coli* occurs in two modes, running and tumbling, which are defined by the direction of the flagellar spin and, ultimately, the signal transduction response that arises from interaction of chemoreceptor with ligand [42]. Bacteria in the presence of an attractant will undergo prolonged running responses with low tumbling frequency [42, 43]. To observe the effects of synthetic ligands on tumbling frequency, *E. coli* were treated with galactose or galactose-bearing ligands, and bacterial motion was recorded and analyzed using the method of Sager *et al.* [44]. The tumbling frequency was assessed by averaging the mean angular velocity of the paths obtained in the first 5-15 seconds after addition of attractant (Figure 2A). When bacteria were treated with increasing concentrations of galactose, the mean angular velocity decreased, indicative of a running response. Figures 2B-E illustrate sample paths for representative bacteria treated with buffer alone, galactose, compound 1 and compound 3. Treatment with monovalent compound 1 produced similar effects to that of the free chemoattractant (galactose), indicating that the anomeric substituent in 1 did not preclude chemotactic activity. Multivalent compound 3 was more active than monovalent 1 or unmodified galactose. Multivalent compound 3 induced a low mean angular velocity even at very low (e.g., 0.001 mM) saccharide residue concentrations. The response of the bacteria to 3 at 0.1 mM saccharide residue concentrations (ca. 0.004 mM concentration) was comparable to that obtained at ten fold higher (1 mM) concentrations of unmodified galactose. The observed differences in concentration of maximum activity between the monomer 1 and multivalent 3 demonstrate that ligand valency affects chemotactic activity.

*E. coli* were subjected to concentration gradients of compounds 1-4 in capillary accumulation assays [45] to determine the concentration at which the maximum chemotactic response is achieved and the number of bacteria that accumulate at this maximum [34].

When compounds 1-4 were used as attractants in the capillary accumulation assay, oligomer 2 was no more active than monovalent 1; both elicited a maximum chemotactic response at 1 mM (Figure 3A). Compound 2 displays a higher local concentration of galactose to the receptor, however, the similarity of activities for 1 and 2 indicates that a high local concentration of attractant does not alone give rise to increased chemotactic activity. For compounds 3 and 4, concentrations of maximum chemotaxis were significantly lower; the maximum for 3 is at a galactose residue concentration of 0.25 mM (ca. 0.01 mM ligand concentration, 100-fold lower than free galactose) and the maximum for 4 is at a galactose residue concentration of 0.10 mM (0.0002 mM ligand concentration. Concentrations of

maximum chemotaxis of 3 and 4 are 100- and 5000-fold lower, respectively, than free galactose (Figure 3B). The ligands of higher valency (3 and 4), therefore, can induce chemotaxis at extremely low concentrations.

Chemotaxis receptors have been found to be approximately 90 Å apart [46].

Molecular modeling studies indicate that the maximum length of oligomer 2 is approximately 50 Å [23]. The significantly higher potency of the longer oligomers 3 and 4, compared to that of oligomer 2, is believed to be due to the ability of the longer oligomers to cluster chemotaxis receptors. Compound 1 was not a chemoattractant for *ggbp* (AW550 and AW543) or *trg* (AW701) *E. coli* mutants. The results obtained indicated that the ligands 1-4 act specifically to affect chemotaxis through the galactose-sensing machinery.

The number of *E. coli* accumulated in assays employing 1-4 (see Figures 3A and 3B) is less than that when galactose is used as an attractant (120,000 bacteria [34]), despite the observed potency of these ligands in the video assays (see Figure 2A). Capillary accumulation assays depend on proper bacterial reorientations to travel into the capillary for collection. The potency of these ligands may disrupt the ability of bacteria to reorient, decreasing the apparent number of bacteria accumulated. At a given saccharide residue concentration of a multivalent ligand, fewer molecules are present to activate the receptors, and these molecules must traverse the outer membrane. These features of the system may also contribute to the decreased numbers of bacteria accumulated.

To test the generality of the observed valency-dependent differences in chemotactic activities and to investigate the role of membrane permeability in responses to our ligands, chemotactic experiments in *B. subtilis* were conducted. *B. subtilis* is a gram-positive bacterium that, like gram-negative *E. coli*, is able to respond to saccharide chemoattractants [47, 48]. In the case of *B. subtilis*, the multivalent ligands can directly interact with saccharide-sensing receptors, without having to first traverse the outer membrane. Glucose is a chemoattractant for *B. subtilis* [47], but galactose is not. Glucose-carrying ligands (compounds 5-7, Scheme 1) were effective chemoattractants for *B. subtilis* as shown in capillary accumulation assays. In addition the chemotactic responses to glucose-carrying ligands were shown to also depend on ligand valency. As shown in Fig. 4, monomer 5 elicited maximum activity at 1 mM, while oligomer 6 elicited maximum activity at a saccharide residue concentration of 0.1 mM (50-fold lower ligand concentration than that of free glucose) and oligomer 7 elicited maximum activity at a saccharide residue concentration of 0.01 mM (1250-fold lower ligand concentration than that of glucose). Free

chemoattractant signal glucose had maximal activity as a chemoattractant at 0.5 mM. In analogy to observations with *E. coli*, as the valency of the ligand increases, the saccharide residue concentration of maximum chemotaxis decreases. Significantly, the number of bacteria accumulated towards 5-7 was comparable to the number accumulated when unmodified glucose was used as the attractant. Consistent with previous reports on the activity of galactose, galactose-bearing ligands (such as 1) were not chemoattractants for *B. subtilis* [47], further indicating that the multivalent ligands were acting specifically. The results observed indicate that in evolutionarily divergent bacteria *E. coli* and *B. subtilis*, the valency of the attractant influences chemotactic response.

Fluorescence microscopy experiments were performed to visualize changes in chemotaxis receptor organization upon treatment with saccharide-carrying ligands. These experiments can determine directly whether or not multivalent ligands can influence chemoreceptor reorganization. It had been shown that wild-type *E. coli* localize chemoreceptors to their poles and that inactivation of the structural protein, CheW, results in a random distribution of chemoreceptors on the cell [49]. The ability of ROMP-derived arrays to localize the chemoreceptors was examined using *E. coli cheW* mutants. Bacteria were treated with 1, 3, or 4, fixed, and labeled with an antibody to the bacterial chemoreceptors (anti-Tsr). Monovalent compound 1 had no effect on receptor distribution, but multivalent compounds 3 and 4 were observed to reorganize the chemoreceptors. As anticipated, localized receptors in the *cheW* cells occurred at seemingly random locations, in contrast to the polar localization observed in the wild type bacteria. Receptor clustering was more pronounced in the case of cells treated with the longer oligomer 4 than with 3. The results of these experiments indicate that ROMP-derived multivalent compounds can induce lateral receptor reorganization. The differences observed in chemotactic activities of the multivalent ligands as a function of oligomer length and the observation that ROMP-derived multivalent ligands can induce lateral receptor reorganization supports the conclusion that receptor reorganization is involved in mediating the chemotaxis response. The results further indicate that changes in receptor localization can give rise to changes in chemotactic responses.

To confirm the ability of multivalent ligands to alter the organization of the chemoreceptors in the bacterial membrane *E. coli* were treated with compound 8, a galactose-bearing multivalent ligand having a fluorescent label (Scheme 1). When *E. coli* were treated with 8 or a fluorescein-labeled anti-Tsr antibody, the fluorescence patterns observed were

AB9  
cont

5 similar. Both materials were observed to bind at the poles of the bacteria indicating that the ROMP-derived ligands bind specifically to the bacterial chemoreceptors. To address directly the ability of these multivalent ligands to reorganize receptors, *CheW* mutants were treated with both compounds. Patches of anti-Tsr antibody labeled chemoreceptors that colocalize with compound 8 were observed, as illustrated in Fig. 6. This result indicates that multivalent ligand 8 is responsible for the observed changes in cell receptor organization.

10 The data obtained indicate that multivalent ligands influence chemotactic responses by altering the organization of cell surface chemoreceptors. An alternative view is that these changes are derived from increases in functional affinity, which result from multivalent presentation. While this mechanism is possible, evidence linking changes in ligand affinities with chemotactic activity is lacking. Equilibrium-binding constants for various ligands often do not correlate with ligand activities in bacterial chemotaxis assays [34, 35, 50, 51]. In contrast, a number of studies have implicated receptor localization in chemotaxis [38-41, 46, 52]. It has been shown, for example, that assembled tetramers of the chemoreceptor Tar are more active in *in vitro* signaling than are individual receptors or dimers [53]. Together, the present data and these results suggest that the differences in chemotactic activities for monovalent 1 versus multivalent 3 and 4 are due to their abilities to control the valency of receptor clusters. Based on these result, a mechanism in which systematic increases in ligand valency lead to changes in chemotactic responses by incorporation of additional receptors into clusters (as illustrated in Figures 7A-D) is proposed.

25 By generating synthetic molecules using ROMP that differ only in ligand valency, as opposed to ligand density or spacing, it has been shown that the valency of a ligand influences its ability to organize chemoreceptors and its ability to elicit a chemotactic response from those receptors. The results demonstrate that multivalent ligands of distinct valency (distinct or defined number of functional moieties), such as those described herein, can be used to tune cellular responses through changes in receptor organization. Further, ligand valency can be used to tune chemotactic responses of diverse bacteria (both *E. coli* and *Bacillus subtilis*) indicating that the methods of this invention are generally applicable to diverse cell types. The ROMP-based synthetic route to multivalent arrays is general [54] and can be employed to generate a variety of multivalent ligands or arrays which carry a variety of types and numbers of chemical signals that bind to cell receptors (cell surface receptors, transmembrane receptors and cytoplasmic receptors) and which as a result, likely mediated by lateral receptor reorganization, elicit a biological response. Control of the type of signal

30

covalently bonded to the multivalent ligand and control of the spacing and number of signals presented on the ligand can be used to tune the type and magnitude of the response elicited.

It has also been found that multivalent ligands that bind to one type of receptor can affect the biological response induced by binding of ligands to another type of receptor.

Serine is another small molecule (in addition to galactose) which acts as a chemoattractant for bacteria, such as *E. coli*. Initial contact of *E. coli* cells with a multivalent ligand with galactose SREs, compounds **2** and **3** was followed, after a 2 min adaptation period, by addition of varying concentrations of serine. The chemoattractant effect of serine was enhanced about 30%, measured as average mean velocity (deg/frame) (see Figs. 5A and B), in the presence of multivalent ligands compared to serine in the absence of the multivalent ligand. It is believed that clustering of galactose-binding cell receptors by the multivalent ligand caused the enhancement of the response of the cell to the other chemoattractant serine, see Fig. 1.

#### Modulation of Neutrophil Chemotaxis

Neutrophil migration is an example of cell migration. Neutrophils migrate toward a number of different endogenous and exogenous substances. N-formyl peptides are one type of exogenous substance that is a chemoattractant for neutrophils [65], a bacterial transcription by product. Neutrophils have cell surface receptors which bind to the chemoattractant and can sense increasing concentration gradients of the chemoattractant. Neutrophils respond to the chemoattractant by migrating toward increased concentrations leading them to the site of infection, for example. In addition, and also in response to such chemoattractants neutrophils release intercellular signals that affect responses in other cells, particularly other immune systems cells. Multivalent ligands of this invention can be used to enhance the response of neutrophils to chemoattractants and enhance immune system clearance of infectious agents. Scheme 2 illustrates an exemplary N-formyl peptide **20** and an exemplary SRE for that N-formyl peptide **21** for use in multivalent ligands that modulate neutrophil migration. These signal groups (SREs) can be covalently or noncovalently bonded to ROMP scaffolds such as those illustrated in Scheme 2 (**22** and **23**). Scheme 3 provides exemplary linkers that can be employed in multivalent ligands carrying N- formyl-peptides.

## Modulation of Immune Processes

The development of an immune response can be modulated via valency-dependent interactions of immune system cells with multivalent ligands of this invention. The recognition of foreign (non-self) epitopes, cells, viruses or viral particles for clearance by the immune system is due in part to cell receptors that recognize the epitopes, cells, viruses or viral particle as foreign. In order for clearance to occur, the foreign signal must be recognized and there must be a B cell or T cell response to the foreign signal. Proper immune responses require activation and subsequent deactivation of B cells and T cells. Receptor clustering on B cells and T cells has been implicated in the production of an immune response.

Multivalent ligands of this invention which have one or more RE or SRE through which the ligand can bind to a B cell, T cell or other immune cell and which carry one or more antigens, epitopes can be employed to modulate the response of the immune cell (enhancing or decreasing immunogenicity of the antigen or epitope). When the epitope or antigen is recognized as foreign (non-self) by the immune cell, cells or immune system in which an immune cell is found, then the multivalent ligand can be used to tolerize the immune cell, cells or immune system to the epitope or antigen. In this case, the epitope or antigen is that of a beneficial or clinical species (cell, particle, nucleic acid) or of a self cell (or tissue) that is incorrectly recognized as foreign (non-self). In contrast, a multivalent ligand of this invention can be used to sensitize or increase the sensitivity of the immune cell, cells or immune system to the foreign epitope or antigen enhance its immunogenicity and enhance the immune response to it. This method would be employed with a foreign epitope or antigen that was not beneficial, e.g., one associated with a pathogen. When the epitope or antigen is recognized as self by the immune cell, cells or immune system in which an immune cell is found, then the multivalent ligand can be used to sensitize the immune cell, cells or immune system to the self epitope or antigen. In this case, the epitope or antigen may be of a non-beneficial self cell or macromolecule, e.g., a cancer cell, or may be a foreign epitope or antigen that is incorrectly recognized as self. In contrast, a multivalent ligand of this invention can be used to tolerize the immune cell, cells or immune system to a self epitope or antigen that is incorrectly recognized as foreign. Methods for tolerization and sensitization are specifically exemplified hereafter.

The C3d complement fragment binds the CR2 receptor (CD21/CD19 complex) on B cells. The expression fusion product of the fusion of the cloned C3d gene fragment and the

5 C-terminal region of hen egg lysozyme gene was able to increase immunogenicity significantly more (1000-fold) than the level achieved with the lysozyme combined with a strong adjuvant [62]. Scheme 4 illustrates an exemplary multivalent ligand containing two different signal groups 30 prepared from the ROMP polymer 29 by selective covalent bonding of the different signals. One of the signals is a hen egg lysozyme (HEL) peptide (specific for the A20 cell line): 103-117 NGMNAWVAWRNRCKG (SEQID NO: 1)[63] and the other is a 16-mer C3d peptide involved in binding to CR2: KNRWEDPGKQLYNVEA (SEQID NO: 2)[62]. This HEL peptide can be attached to the polymer backbone at the N-terminal amine (40) of the peptide or at a side group of a lysine near the end of the peptide (41):

40: \*GDGNGMNAWVAWRNR-CONH<sub>2</sub> (SEQID NO: 3) or

41: DGNGMNAWVAWRNRGK\*-CONH<sub>2</sub> (SEQID NO: 4)

where \* indicates the site of attachment. The C3d peptide can be attached to the multivalent ligand via the thiol of cysteine positioned at either end of the peptide(42 and 43):

42: \*CKNRWEDPGKQLYNVEA (SEQ ID NO: 5) or

43: KNRWEDPGKQLYNVEAC\* (SEQ ID NO: 6)

Multivalent ligands containing signals 41 alone or in combination with 42 or 43 or 40 alone or in combination with 42 or 43 can induce an enhanced immune response compared to HEL its self. A multivalent ligand containing a plurality of peptide elements that are ligands for the CR2 receptor can cluster the CR2 receptor on the surface of the B cell and as demonstrated in the chemotaxis experiments can enhance the response of that B cell to other ligands, e.g., anitgens. Multivalent ligands containing one or more bound CR2 ligands in combination with one or more bound anitgens can cluster the CR2 receptor with the receptor that recognizes the antigen and thereby enhance the response of the B cell to the antigen.

25 Clustering of CR2 with a receptor that recognized HEL(for example) on the B cell surface can enhance the response of the B cell for the HEL antigen and can result in an enhancement of immune response toward the HEL epitope. An alternative hen egg lysozyme peptide that can be employed in construction of multivalent ligands of this type is:

44: ELAAAMKRHGLDNYRGYSLGNWVCA (SEQID NO: 7).



CD22 is a B cell surface glycoprotein involved in cell adhesion and activation [64]. CD22 is important in the negative regulation of B cell antigen receptor signaling [74]. The structure recognized by CD22 is Sial2 $\alpha$ 6 Gal $\beta$ 14GlcNAc $\beta$  (Scheme 5, compound 50). This signal can be attached to a ROMP polymer backbone as illustrated in Scheme 5 via a primary thiol group (compound 51). Multivalent ligands containing one or more ligands for CD22 (such as 51) in combination with one or more HEL epitopes (such as 42 or 43) attenuates the immune response to the HEL epitope.

#### Crosslinking (Aggregation) of Cells.

Many proteins, such as lectins and antibodies, possess multiple ligand binding sites. When these proteins bind to ligands immobilized on adjacent cell surfaces, the cells aggregate. Cell aggregation can be monitored easily, and this property has found use in the development of diagnostics for pathogen detection [75], therapeutics [76-78], blood typing tests [79], and other biotechnological applications [80-82]. Many lectins have been shown to have mitogenic activities that are dependent on the valency of the lectin. These mitogenic lectins, including ConcanavalinA (ConA), are thought to cluster glycoproteins on the surface of the target cell, activating mitogenic signals and inducing cell proliferation [67, 68]. Lectins have been useful tools for exploring signal transduction [69, 70] and cell growth [71, 72], and studies using them have elucidated possible functional roles for mammalian lectins, such as the galectins and selectins.

The effectiveness of multivalent proteins at instigating cell aggregation is determined by how tightly the protein binds to cell surface ligands. One effective way to increase the avidity of these interactions is to increase the number of ligand binding sites [83-85]. Research efforts have focused on favoring oligomer formation for lectins [86-87] or generating novel multimers of antibody scFv fragments [88]. Methods which further enhance the number of binding sites or favor the optimized orientation of these binding sites would increase the utility of these materials in many applications.

Lectins are a large class of saccharide-binding proteins, many of which are homo-oligomers assembled from two to four copies of identical subunits [89]. Lectins aggregate cells when they crosslink glycoproteins or glycolipids on adjacent cell surfaces. Aggregation can be modulated by altering the number of active monomers within the lectin oligomer. For example, the ability of the tetravalent mannose-binding plant lectin concanavalin A (Con A) to aggregate red blood cells is greatly decreased when the lectin is forced into a lower valency

09815296 " 073001  
5 dimeric form by succinylation [87]. Increasing the valency of lectins may have the opposite effect, i.e. to enhance cell aggregation; however, methods have not been readily available for generating lectin complexes with higher order valencies. Because the valency of ROMP-derived materials can be altered systematically, the effect that the number of saccharide groups, such as mannose, bound to the ligand has on the number of lectins, such as ConA, assembled on a given scaffold can be investigated.

10 The precipitation of Con A depends on the clustering of Con A tetramers and this technique can be used to determine the stoichiometry of insoluble Con A - ligand complexes [90]. To investigate the formation of Con A clusters with multivalent ligands of this invention, ROMP-based scaffolds containing defined numbers of mannose residues, the monomer **9** and polymers **10-13** having n of 10, 25, 50, or 100, respectively, illustrated in Scheme 1 were prepared using ROMP methods [54]. Compounds **9-13** were contacted with Con A, monomeric compound **9** was unable to induce precipitation, but multivalent compounds **10-13**, caused concentration-dependent precipitation of Con A. Precipitation results further indicated that the stoichiometry of ConA complexed with **10** (the 10-mer) is about 2:1 and that of complexes of ConA with **11** and **12** is approximately 4:1.

20 In contrast, dimeric succinylated Con A precipitated only with the highest valency multivalent ligand compounds tested, e.g., compound **12**, and the complexes formed had a 4:1 (receptor:scaffold) stoichiometry in the precipitate. Thus, the number of mannose residues displayed by the scaffold is important in the formation of protein-scaffold complexes. Precipitation results were substantially confirmed with a transmission electron microscopy (TEM) technique in which clusters of biotinylated ConA with compounds **10-12** were labeled with a high density streptavidin – gold particle. Compound **10** was observed to form dimers exclusively, while **11** was able to form both dimers and trimers and compound **12** formed both dimers and trimers as well, but favored trimeric clusters more than the other scaffolds.

25 The assembly of Con A clusters in solution can be monitored by fluorescence resonance energy transfer (FRET), in which fluorescein and tetramethylrhodamine (TMR) serve as donor and acceptor fluorophores [91, 92]. When these fluorophores are within approximately 80 Å the fluorescein signal is quenched, such that fluorescein fluorescence should decrease when labeled Con A is assembled into clusters [93]. Compounds **9-12** were added to a solution of fluorescein- and TMR- labeled Con A. The fluorescence emission maximum of fluorescein was monitored to ascertain which scaffolds promoted the formation

of Con A clusters. In agreement with the previous experiments, Con A clusters formed in the presence of multivalent ligands 10-12 but not with monomeric compound 9. The fluorescence quenching was dependent not only on scaffold valency, but also on ligand concentration. Quenching first increased as scaffold concentration increased and then decreased again as the concentration was increased further. The absence of quenching at high scaffold (multivalent ligand) concentrations indicates that Con A clusters are disfavored at these concentrations, likely because of site saturation. The high concentration of scaffold compared to Con A favors occupation of each ligand binding site on Con A by individual polymers precluding clustering of multiple lectins.

The ability of Con A clusters formed on ROMP-derived polymers to aggregate Jurkat cells was examined initially by light microscopy (see Fig. 10). Con A alone was able to induce some Jurkat cell aggregation even at low concentrations (5  $\mu\text{g/mL}$ ). When monovalent Con A ligands such as methyl  $\alpha$ -D-mannopyranoside or 9 were premixed with Con A they inhibited aggregation, presumably by destabilizing Con A - cell interactions. For Jurkat cells, inhibition occurred even at low concentrations (0.5  $\mu\text{M}$ ) of monovalent ligands. Interestingly, multivalent compounds 10-12 did not inhibit Jurkat cell aggregation at 0.5  $\mu\text{M}$ , a concentration shown to be optimum for Con A cluster formation under similar conditions. Increasing the concentration of the multivalent ligand 10-fold (5  $\mu\text{M}$ ) abolished aggregation activity, consistent with site saturation. Thus it is possible to alternatively inhibit or promote cell surface-lectin interactions by varying scaffold valency and multivalent ligand concentration. The ability of Con A complexed to multivalent ligands to interact with cell surfaces was thus tunable.

Further experiments were conducted which demonstrated that ConA-mediated agglutination of erythrocytes could be controlled by addition of multivalent ligands (compounds 9-13). Certain combinations of ConA and multivalent ligands exhibited enhanced agglutination of these cells compared to ConA itself, as shown in Fig. 11. In particular, a combination of ConA tetramer and multivalent ligand (compound 13) at concentration ratio 10:1 (based on tetrameric ConA and based on the number of mannose residues) exhibited significantly enhanced agglutination compared to ConA alone.

Complexes containing multiple Con A tetramers were assembled readily on compounds 10-13 when intermediate multivalent ligand concentrations were used, but were not detectable when the concentration of the scaffold was either too low or too high. The

1144  
concentration range over which such complexes are formed depends upon the relative concentrations of ConA and multivalent ligand (based on the number of ligands, RE or SRE) and upon the valency of multivalent ligand. This is generally true for any complex of a multivalent ligand with any protein. The concentration range over which complexes of a multivalent ligand with one or more ConA (or such complexes with any lectin or more generally with any protein) can be readily determined for a particular application under particular conditions by assessing retention of function by ConA (or more generally the protein or lectin). Complexes of multivalent ligands with ConA will generally be formed, dependent upon the valency of the multivalent ligand and the particular experimental conditions, when the concentration range of the ligand (based on numbers of SRE, e.g., mannose) ranges from about 1:1 to over 100:1.

09015296.073001  
The results herein indicate generally that the valency and concentration of a multivalent ligand can be varied to control the assembly of lectin on to the multivalent scaffolds of these multivalent ligands. More specifically, the valency and concentration of ROMP-derived materials can be varied to control the formation of Con A clusters, as illustrated in Figs. 9A-C. Monovalent ligands (as well as low concentrations of multivalent ligands) bind to lectin, but do not inhibit cell aggregation (Fig. 9A). Under conditions that favor lectin-scaffold complexation, i.e., intermediate concentration levels of multivalent ligands, a plurality of lectins can be assembled on the multivalent ligand and the lectins retain free saccharide binding sites capable of interacting with cell surfaces (Fig. 9B). When multivalent ligand concentration is increased, lectin binding sites are saturated by binding to a plurality of multivalent ligands, lectin assembly is disfavored and lectins are not capable of interacting with cell surfaces (Fig. 9C). Thus, as illustrated, scaffold valency and ligand concentration can be controlled to assemble lectin clusters with multivalent ligands wherein the lectin retains cell binding activity. Further, scaffold valency and more importantly multivalent ligand concentration can be controlled to inhibit the cell aggregation function of lectins.

These results demonstrate that proteins, such as lectins, can be assembled on a polymeric scaffold, such as those provided by the multivalent ligands of this invention, and that the assembled proteins, including lectins, will retain biological function. Methods described herein can be employed to generate polymeric assemblies of one or more lectins, as well as polymeric assemblies of one or more antibodies or antibodies fragments, which retain the ability to bind to ligands (e.g., saccharides or epitopes). Methods herein are generally

applicable to generation of assemblies of various chemical and biological species, particularly macromolecular species, including proteins, carbohydrates, nucleic acids though binding to recognition elements and signal recognition elements in a multivalent ligand.

#### Enhancement of Cell Toxicity Using Multivalent Ligands

Lectins, such as Con A, as well as agglutinins and phytohemagglutinins in general, can exhibit toxic effects in certain kinds of cells. Multivalent ligands carrying saccharide groups can complex with lectins, such as Con A, as discussed above. Complexes containing several lectin molecules complexed to an appropriately substituted multivalent ligand can function to aggregate cells, if binding sites on the lectin are not saturated by binding to the ligand groups. When higher multivalent ligand concentrations (dependent upon the specific conditions and applications, and dependent upon ligand valency) are employed, lectin binding sites can become saturated and cell aggregation by the lectin is then inhibited. Saturation of a given lectin by a given multivalent ligand can be readily determined empirically. Further, saturation of the function of any protein by a given multivalent ligand can be determined by assessing function of the complexed protein.

Complexes of a lectin with multivalent ligands have been found to exhibit cell toxicity that is enhanced over that of the lectin itself. As illustrated in Fig. 12, PC12 cells treated with 0.1  $\mu$ M Con A (for 48 hr) exhibited no apparent loss of viability. In contrast, PC12 cells treated with combination of 0.1  $\mu$ M Con A and 4  $\mu$ M of compound **11** under the same conditions exhibit almost a 30% loss in viability. These results indicate that complexes of lectin with multivalent ligands of this invention in which the ratio of the concentrations of ligand to lectin is sufficiently high to saturate ligand binding sites of the lectin can trigger apoptosis in cells.

#### Generation of Multivalent Polymers

ROMP was used to convert **1** to the series of oligomers **2-4** as previously described [55]. Similar conditions were employed in the synthesis of oligomers **6** and **7** [54]. Fluorescent polymer **8** was generated by specific end-labeling with a bifunctional capping agent [Scheme 7] and subsequent conjugation to the fluorophore BODIPY-TR (commercially available from Molecular Probes) [56]. Compounds **9-12** were the samples prepared and tested in reference [54]. The degree of polymerization (dp) for each compound was determined by  $^1\text{H}$  NMR.

Valency (n) is an approximation of the degree of polymerization (DP), where DP is the ratio of monomer to catalyst used in ROMP.

### Video Microscopy

*E. coli* AW405, which exhibits wild-type chemotactic responses, from an overnight culture were grown in LB (Luria Bertani broth) to OD<sub>550</sub> of 0.4-0.6 and then washed twice with attractant-free chemotaxis buffer (10 mM potassium phosphate buffer, pH 7.0, 10  $\mu$ M EDTA). Partially permeabilized bacteria (25  $\mu$ M EDTA for 3 min. at room temperature, then quench with 50  $\mu$ M CaCl<sub>2</sub>) at an OD<sub>550</sub> of 0.1 were placed under a cover slip supported by additional cover slips in the method of Sager *et al.* [44]. (Permeabilization had no effect on bacterial chemotaxis toward galactose or 1 but was necessary for chemotaxis toward 4 [57]). Bacteria were allowed to adjust to contact with glass surface for 1-2 min. Attractant was added to achieve the final concentration indicated at a 5  $\mu$ L final volume. The bacterial motion at 28 °C was recorded, and the paths were analyzed using the ExpertVision system. Paths derived from the first 5 to 15 seconds following the introduction of attractant were analyzed. Angular mean velocities varied approximately 14% between experiments performed on different days. Data were analyzed using the Q and Students tests.

### Capillary Accumulation Assay

*E. coli* from an overnight culture were grown in LB to OD<sub>550</sub> 0.4-0.6, washed twice with *E. coli* chemotaxis buffer, and then partially permeabilized. Bacteria were resuspended in chemotaxis buffer to an OD<sub>550</sub> 0.1 and utilized in the capillary accumulation assay at 30 °C for 60 min, as previously described [45]. *B. subtilis* OI1085 was grown from an overnight culture in T broth (1% tryptone, 0.2 mM MgCl<sub>2</sub>, 0.5% NaCl, 0.01 mM MnCl<sub>2</sub>) supplemented with 10 mM glucose and 0.5% glycerol, washed with *B. subtilis* chemotaxis buffer (10 mM phosphate buffer, pH 7.0, 10  $\mu$ M EDTA, 0.5% glycerol, 0.3 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>), and capillary assays were performed at a final OD<sub>550</sub> 0.01 at 37 °C for 30 min [47]. The number of *B. subtilis* accumulated was normalized to 500 bacteria accumulated towards buffer alone. Results of capillary assays can be influenced by factors other than the activity of the attractant, such as metabolism of the substrate or toxicity [45, 58]. To exclude this possibility, we tested the ability of *E. coli* to utilize 1 as a sole carbon source. These experiments revealed that 1-4 are not toxic and that monomer 1 is not metabolized (data not shown). Data was analyzed using the Q and Students tests.

## Immunofluorescence Microscopy

*E. coli* AW405 or RP1078 (*cheW*) were pretreated with buffer alone or with compounds 1, 3, 4, or 8 at 5 mM in a 10  $\mu$ L total volume of chemotaxis buffer. After a 10 minute incubation at 30 °C, the bacteria were fixed (2% paraformaldehyde (PFA) in HEPES pH 7.0, 30 min., 4 °C), placed on poly-L-lysine treated cover slips in the bottom of 6-well plates, permeabilized with methanol, and labeled with anti-Tsr antibody (1:250) and fluorescein-labeled goat-anti-rabbit antibody (1:500) according to the procedure of Maddock and Shapiro [49]. Anti-Tsr antibodies recognize the conserved chemoreceptor cytoplasmic domain and are thus cross-reactive with multiple chemoreceptors. Some binding exclusion (exclusive 530 nm or 590 nm fluorescence at a pole) was seen in dual labeling experiments in which both antibody and 8 were used. Fluorescence microscopy was performed on a Zeiss Axioscope at 1000x using an oil immersion lens. Images were captured using IPLab Spectra 3.2 and Adobe PhotoShop 5.0.

## Quantitative Precipitation

Quantitative precipitations and analysis were carried out by a method modified from that previously described by Khan, *et al* [90]. Briefly, Con A (Vector Laboratories, Burlingame, CA) and scaffold were dissolved in precipitation buffer (0.1 M Tris-HCl pH 7.5, 90  $\mu$ M NaCl, 1 mM  $\text{CaCl}_2$ , 1 mM  $\text{MnCl}_2$ ), vortexed briefly to mix, and then incubated for 5 hours at room temperature (or 2 days at 4 °C for succinylated Con A). The final concentration of Con A tetramers was 30  $\mu$ M (assuming Con A tetramers with a molecular weight of 104,000) and succinylated Con A dimers was 44  $\mu$ M (assuming dimers with a mass of 52,000). White precipitates were pelleted by centrifugation at 5000 xg for 2 minutes. Supernatants were removed by pipet and pellets were gently washed twice with cold buffer. Pellets were then resuspended in 600  $\mu$ L 100 mM methyl  $\alpha$ -D-mannopyranoside (100  $\mu$ L for succinylated Con A), and were completely dissolved after a 10 minute incubation at room temperature. Protein content was determined by measuring the absorbance at 280 nm by UV-vis spectroscopy on a Varian Cary 50 Bio using a 100  $\mu$ L volume quartz cuvette. Measurements are the average of three independent experiments.

## Transmission Electron Microscopy

TEM methods were performed essentially as previously described [96]. Con A tetramers were labeled with biotin using conditions that favored attachment of 1-2 copies of

biotin residues. Biotinylated ConA tetramers were mixed with ligands of interest in solution and then contacted with an excess of streptavidin-conjugated 10 nm gold particles. Samples can be treated with 2% phosphotungstic acid (pH 7.0, 30 sec) to enhance contrast. Images of random fields were acquired for each treatment and analyzed for formation of ConA  
5 complexes. Gold particles within 25 nm of less of each other were considered to be part of a complex. This distance was based on the modeled length of the synthetic multivalent ligands used [23] and the structure of tetrameric ConA determined by X-ray crystallographic analysis[97].

Specifically, biotinylated Con A (2.3  $\mu$ M) and scaffold (0.75  $\mu$ M, mannose  
10 concentration) in PBS pH 7.2 were incubated for 15 minutes at room temperature before streptavidin - 10 nm gold (Sigma, St. Louis, MO) was added to a final concentration of 3.0  $\mu$ M. Complexes were incubated at room temperature for 15 minutes and then placed onto carbon-coated Formvar-treated grids. Grids were air dried and viewed on a LEO Omega 912 Energy Filtering Electron Microscope (EFTEM). Images were at 12,500 x magnification,  
5 collected on a ProScan Slow Scan CCD camera, and analyzed in Adobe PhotoShop 5.0. Fields averaged between 5 and 50 gold particles and 15-20 fields were collected on each day for each treatment. Results are the average of results obtained on three separate experiments performed on independent samples on three separate days. Total number of gold particles collected on each day varied from about 80 to over 400.

#### Fluorescence Resonance Energy Transfer

Fluorescein-Con A (Vector Laboratories, Burlingame, CA) and TMR-Con A (Sigma,  
St. Louis, MO) in phosphate buffered saline (PBS) pH 7.2 were mixed to afford final  
concentrations of 4  $\mu$ g/mL and 0.4  $\mu$ g/mL respectively. Scaffold was added in PBS to the  
25 final concentrations indicated, with a final volume of 200  $\mu$ L. This solution was vortexed briefly and then incubated at room temperature for at least 15 minutes. No precipitates were observed in any of these samples. Fluorescence was determined on a Hitachi F-4500  
fluorospectrophotometer using a 200  $\mu$ L volume quartz cuvette, an excitation wavelength of 492 nm, emission wavelength of 515 nm, and 10 nm slit widths. Emission intensities are the  
30 average of 3-5 independent experiments with 3 scans performed during each experiment. Compounds 9-12 had negligible fluorescence at 515 nm. Curves were fit using the equation:

$$\%F = (\%F_{\max} \times L)/(L + IC_{50})$$



where %F is the change in fluorescence relative to untreated,  $\%F_{\max}$  is the maximal recovery of fluorescence, L is the micromolar mannose concentration of scaffold, and  $IC_{50}$  is the half-maximal concentration for inhibition of clustering.

#### Jurkat Cell Aggregation

Jurkat cells were cultured and maintained as previously described [94]. Cells were washed three times in cold PBS and then treated with Hoescht 33342 (100  $\mu$ g/mL) for 30 minutes at 30 °C. Cells were washed twice with cold PBS and then fixed for 30 minutes at 4 °C with 2% paraformaldehyde in HEPES pH 7.4. Fixed cells were washed twice and then treated in 200  $\mu$ L final volume with premixed solutions of Con A and scaffold. A 2x solution of Con A and scaffold was prepared in PBS pH 7.2, vortexed briefly, and then incubated at 22 °C for 30 minutes before being added to cells. Cells, Con A solutions, and 100  $\mu$ g/mL DNase (to prevent cell aggregation by nucleic acid) were incubated at 22 °C for 30 minutes. Cells were pelleted at 400 xg, resuspended gently into 50  $\mu$ L PBS, and then added to slides for visualization at 200 x magnification on a Zeiss Axioscope outfitted with the appropriate filter set. Approximately 100 - 200 cells were counted from random fields on each day. Clusters were scored for at least two cells in direct contact with each other and expressed as a percentage of the total number of objects (individual cells and clusters) counted. Results are summarized in Fig. 10. ROMP-derived ligands 9-12 alone were not able to cause cell aggregation. Images were captured in IPlab Spectra 3.2 and prepared in Adobe Photoshop 5.0.

#### Erythrocyte Agglutination

ConA (53 nM, 5  $\mu$ g/mL) and ligand compound 13 (530 nM; per saccharide basis) were added in an end final volume of 100  $\mu$ L, PBS pH 7.2 in a 96-well plate. The complexes were incubated for 15 minutes at room temperature

#### Cell Toxicity Experiments

HBS buffer contained HEPES (10mM), NaCl (150mM), and  $CaCl_2$  (1mM) at pH = 7.4. Concanavalin A (ConA) was obtained from Vector Labs (Burlington, CA) and was freshly diluted for all experiments. The concentration of the ConA stock solution was determined using  $A_{280}^{1\%} = 13.7$  (95). A single ConA dilution was then made and split for each sample. Ligands were then added from appropriate stock solutions at 5 times the

desired final concentration. All samples had six replicates for each concentration. Control samples were used in each run that contained HBS alone, ConA in HBS and the highest concentration of ligand in HBS. Lysis controls were made by adding HBS buffer alone and adding lysis buffer after 48 h sample incubation.

**Cell Culture:** All cell culture reagents were obtained from GIBCO BRL unless otherwise noted. PC12 cells (ATCC: CRL-1721) were grown in media containing 84% (v/v) RPMI 1640 (with L-glutamine), 5% (v/v) heat inactivated fetal bovine serum, 10% (v/v) heat inactivated horse serum, and 1% penicillin/streptomycin (10000 units/ml), in a humidified incubator at 37°C and 5% CO<sub>2</sub>. Low serum media contained 97.5%(v/v) RPMI 1640 (with L-glutamine), 0.5% inactivated fetal bovine serum, 1% (v/v) heat inactivated horse serum, and 1% penicillin /streptomycin (10000 units/ml). Cells were grown in T-flasks treated with collagen, and harvested at confluence by trypsinization (0.05% trypsin and 0.4 mM EDTA) followed by quenching with fresh medium. Cells were concentrated to pellet (2100 rpm for 10 min), aspirated then resuspended in fresh medium. The population was determined by haemocytometer and treatment with trypan blue, cells were then plated to 96-well plates (tissue culture treated obtained from CoStar, Corning NY) at ~15,000 cells/well. Plates were then incubated for 34 h to allow cells to adhere.

The medium was then removed and replaced with low serum medium(80 µL). Samples and controls in HBS (20 µL) were then added and incubated for 48 h at 37 C. After incubation 10 µL of a 5mg/mL solution of MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (Sigma/Aldrich, Milwaukee, WI) in low serum RPMI medium was added to each well.

After 4 h, 100 µL of lysis buffer (50% dimethyl formamide/ 20% sodium dodecyl sulfate in HBS, pH = 4.7) was added and the cells were incubated overnight. The plate was then read on a plate reader (Biostar) at 570 nm. Percent cell viability was determined using

the following equation:  $\frac{G_1 - G_0}{G_{con} - G_0} = \%V$ , where  $G_0$  is the lysed cell control,  $G_{con}$  is

control cells treated only with vehicle, and  $G_1$  is a sample treated with peptide and vehicle. Results from an experiment in which cells are initially treated with multivalent polymer

compound 11 followed by treatment with ConA and appropriate controls are illustrated in Fig. 11.

5  
10  
Those of ordinary skill in the art will appreciate in view of the descriptions herein that there are a variety of alternative structures, methods, procedure and techniques that can be readily applied or adapted to the practice of this invention other than those that have been specifically exemplified. It will be appreciated that there are a wide variety of designs for and methods for preparation of multivalent ligands with properties as described herein. It will also be appreciated that there are a wide variety of molecular scaffold available for the productive presentation of SRE as well as a wide variety of SRE that can be applied or adapted to the methods described herein.

## References:

1. Mammen, M., Choi, S.-K. & Whitesides, G. M. (1998). Polyvalent interactions in biological systems: implications for design and use of multivalent ligands and inhibitors. *Angew. Chem., Int. Ed. Engl.* **37**, 2755-2794.
2. Kiessling, L. L. & Pohl, N. L. (1996). Strength in numbers: Non-natural polyvalent carbohydrate derivatives. *Chem. Biol.* **3**, 71-77.
3. Williams, L. T. (1989). Signal transduction by the platelet-derived growth factor receptor. *Science* **243**, 1564-1570.
4. Heldin, C.-H. (1995). Dimerization of cell surface receptors in signal transduction. *Cell* **80**, 213-223.
5. Fire, E., Brown, C. M., Roth, M. G., Henis, Y. I. & Petersen, N. O. (1997). Partitioning of proteins into plasma membrane microdomains: clustering of mutant influenza virus hemagglutinins into coated pits depends on the strength of the internalization signal. *J. Biol. Chem.* **272**, 29538-29545.
6. Germain, R. N. (1997). T-cell signaling: The importance of receptor clustering. *Curr. Biol.* **7**, R640-644.
7. Holowka, D. & Baird, B. (1996). Antigen-mediated IgE receptor aggregation and signaling: A window on cell surface structure and dynamics. *Annu. Rev. Biophys. Biomol. Struct.* **25**, 79-112.
8. Monks, C. R. F., Freiberg, B. A., Kupfer, H., Sciaky, N. & Kupfer, A. (1998). Three-dimensional segregation of supramolecular activation clusters in T-cells. *Nature* **395**, 82-86.
9. Yap, A. S., Briehner, W. M., Pruschy, M. & Gumbiner, B. M. (1997). Lateral clustering of the adhesive ectodomain: a fundamental determinant of cadherin function. *Curr. Biol.* **7**, 308-315.
10. Hato, T., Pampori, N. & Shattil, S. J. (1998). Complementary roles for receptor clustering and conformational change in the adhesive and signaling functions of integrin  $\alpha_{IIb}\beta_3$ . *J. Cell Biol.* **141**, 1685-1695.
11. Kuduk, S. D., Schwarz, J. B., Chen, X.-T., et al. (1998). Synthetic and immunological studies on clustered modes of mucin-related Tn and TF O-linked antigens: The preparation of a glycopeptide-based vaccine for clinical trials against prostate cancer. *J. Am. Chem. Soc.* **120**, 12474-12485.



25. Arimoto, H., Nishimura, K., Kinumi, T., Hayakawa, I. & Uemura, D. (1999). Multivalent polymer of vancomycin: Enhanced antibacterial activity against VRE. *Chem. Comm.* **15**, 1361-1362.
26. Gibson, V. C., Marshall, E. L., North, M., Robson, D. A. & Williams, P. J. (1997). Thymine functionalized polymers *via* living ring-opening metathesis polymerisation. *J. Chem. Soc., Chem Commun.* 1095-1096.
27. Sanders, W. J., Katsumoto, T. R., Bertozzi, C. R., Rosen, S. D. & Kiessling, L. L. (1996). L-selectin-carbohydrate interactions: Relevant modifications of the Lewis x trisaccharide. *Biochemistry* **35**, 14862-14867.
28. Gordon, E. J., Sanders, W. J. & Kiessling, L. L. (1998). Synthetic ligands point to cell surface strategies. *Nature* **392**, 30-31.
29. Parkinson, J. S. (1993). Signal transduction schemes of bacteria. *Cell* **73**, 857-871.
30. Hazelbauer, G. L., Berg, H. C. & Matsumura, P. (1993). Bacterial motility and signal transduction. *Cell* **73**, 15-22.
31. Alon, U., Surette, M. G., Barkai, N. & Leibler, S. (1998). Robustness in bacterial chemotaxis. *Nature* **393**, 18-19.
32. Barkal, N. & Leibler, S. (1997). Robustness in simple biochemical networks. *Nature* **387**, 913-917.
33. Grebe, T. W. & Stock, J. (1998). Bacterial chemotaxis: The five sensors of a bacterium. *Curr. Biol.* **8**, R154-R157.
34. Adler, J., Hazelbauer, G. L. & Dahl, M. M. (1973). Chemotaxis towards sugars in *Escherichia coli*. *J. Bacteriol.* **115**, 824-847.
35. Hazelbauer, G. L. & Adler, J. (1971). Role of the galactose binding protein in chemotaxis of *Escherichia coli* toward galactose. *Nature New Biol.* **230**, 101-104.
36. Silversmith, R. E. & Bourret, R. B. (1999). Throwing the switch in bacterial chemotaxis. *Trends Microbiol.* **7**, 16-22.
37. Mesibov, R., Ordal, G. W. & Adler, J. (1973). The range of attractant concentrations for bacterial chemotaxis and the threshold and size over this range. *J. Gen. Physiol.* **62**, 203-223.
38. Jasuja, R., Keyoung, J., Reid, G. P., Trentham, D. R. & Khan, S. (1999). Chemotactic responses of *Escherichia coli* to small jumps of photoreleased L-aspartate. *Biophys. J.* **76**, 1706-1719.

39. Bray, D., Levin, M. D. & Morton-Firth, C. J. (1998). Receptor clustering as a cellular mechanism to control sensitivity. *Nature* **393**, 85-88.
40. Levit, M. N., Liu, Y. & Stock, J. B. (1998). Stimulus response coupling in bacterial chemotaxis: receptor dimers in signaling arrays. *Mol. Microbiol.* **30**, 459-466.
- 5 41. Duke, T. A. J. & Bray, D. (1999). Heightened sensitivity of a lattice of membrane receptors. *Proc. Natl. Acad. Sci. USA* **96**, 10104-10108.
42. Berg, H. C. & Brown, D. A. (1972). Chemotaxis in *Escherichia coli* analysed by three dimensional tracking. *Nature* **239**, 500-504.
43. Amsler, C. D. (1996). Use of computer-assisted motion analysis for quantitative measurements of swimming behavior in petrichously flagellated bacteria. *Anal. Biochem.* **235**, 20-25.
- 10 44. Sager, B. M., Sekelsky, J. J., Matsumura, P. & Adler, J. (1988). Use of a computer to assay motility in bacteria. *Anal. Biochem.* **173**, 271-277.
45. Adler, J. (1973). A method for measuring chemotaxis and use of the method to determine optimum conditions for chemotaxis by *Escherichia coli*. *J. Gen. Microbiol.* **74**, 77-91.
46. Barnakov, A. N., Downing, K. H. & Hazelbauer, G. L. (1994). Studies of the structural organization of a bacterial chemoreceptor by electron microscopy. *J. Struct. Biol.* **112**, 117-124.
47. Ordal, G. W., Villani, D. P. & Rosendahl, M. S. (1979). Chemotaxis towards sugars by *Bacillus subtilis*. *J. Gen. Microbiol.* **115**, 167-172.
48. Ordal, G. W. (1985). Bacterial chemotaxis: Biochemistry of behavior in a single cell. *Crit. Rev. Microbiol.* **12**, 95-130.
49. Maddock, J. R. & Shapiro, L. (1993). Polar location of the chemoreceptor complex in the *Escherichia coli* cell. *Science* **259**, 1717-1723.
- 25 50. Boos, W. (1969). The galactose binding protein and its relationship to the  $\beta$ -methylgalactoside permease from *Escherichia coli*. *Eur. J. Biochem.* **10**, 66-73.
51. Yaghamai, R. & Hazelbauer, G. L. (1993). Strategies for differential sensory responses mediated through the same transmembrane receptor. *EMBO J.* **12**, 1897-1905.
- 30 52. Li, G. & Weis, R. M. (2000). Covalent modification regulates ligand binding to receptor complexes in the chemosensory system of *Escherichia coli*. *Cell* **100**, 357-365.
53. Cochran, A. G. & Kim, P. S. (1996). Imitation of *Escherichia coli* aspartate receptor signaling in engineered dimers of the cytoplasmic domain. *Science* **271**, 1113-1116.

54. Strong, L. E. & Kiessling, L. L. (1999). A general synthetic route to defined, biologically active multivalent arrays. *J. Am. Chem. Soc.* **121**, 6193-6196.
55. Manning, D. D., Strong, L. E., Hu, X., Beck, P. J. & Kiessling, L. L. (1997). Neoglycopolymer inhibitors of the selectins. *Tetrahedron* **53**, 11937-11952.
- 5 56. Gordon, E. J., Gestwicki, J. E., Strong, L. E. & Kiessling, L. L. (2000). A bifunctional capping agent for ROMP: synthesis and cell binding of a fluorescent neoglycopolymer. *Chem. Biol.* **7**, 9-16.
57. Leive, L. & Kollin, V. (1967). Controlling EDTA treatment to produce permeable *Escherichia coli* with normal metabolic processes. *Biochem. Biophys. Res. Comm.* **28**, 229-236.
- 10 58. Adler, J. (1966). Chemotaxis in bacteria. *Science* **153**, 708-716.
59. Stoddard, B. L. & Koshland, D. E. J. (1992). Prediction of the structure of a receptor-protein complex using a binary docking method. *Nature* **358**, 774-776.
60. Zhang, Y., Gardina, P. J., Kuebler, A. S., et al. (1999). Model of maltose-binding protein/chemoreceptor complex supports intrasubunit signaling mechanism. *Proc. Natl. Acad. Sci. USA* **96**, 939-944.
61. Moudgil, K., Sekiguchi, D., Kim, S., & Sercarz, E. (1997) Immunodominance is independent of structural constraints, *J. Immunology* **159** (6), 2574-2579.
62. Lou, D., & Kohler, H. (1998) Enhanced molecular mimicry of CEA using photoaffinity crosslinked C3d peptide. *Nature Biotechnology* , **16**, 458-462.
63. Gapin, L., et al. (1998) Antigen presentation by dendritic cells focuses T cell responses against immunodominant peptides: studies in the hen egg-white lysozyme (HEL) model, *J. Immunology*, **160**, 1556-1564.
64. Powell, L., & Varki, A. (1994) The oligosaccharide binding specificities of CD22 $\beta$ , a sialic acid-specific lectin of B cells, *J. Biol. Chem.*, **269**, 10628-10636.
- 25 65. Ye, R., & Boulay, F., (1997) Leukocyte Chemoattractant Receptors, *Advances in Pharmacology* **39**:221.
66. PCT application WO 98/46270, (Advanced Medicine) published October 22, 1998.
67. Ramaschi, G., Torti, M., Sinigaglia, F. & Balduini, C. (1993). Intracellular Calcium Mobilization is Triggered by Clustering of Membrane Glycoproteins in Concanavalin A-stimulated Platelets. *Cell Biochem. Function* **11**, 241-249.
- 30 68. Kaplan, M.R., Trubniykov, E. & Berke, G. (1997). Fluorescence Depolarization as an Early Measure of T Lymphocyte Stimulation. *J. Immunol. Methods* **201**, 15-24.



69. Walzel, H., Blach, M., Hirabayashi, J., Kasai, K.-I. & Brock, J. (2000). Involvement of CD2 and CD3 in Galectin-1 Induced Signaling in Human Jurkat T-cells. *Glycobiology* **10**, 131-140.
70. Wenzel-Seifert, K., Krautwurst, D., Lentzen, H. & Siefert, R. (1996). Concanavalin A and Mistletoe Lectin I Differentially Activate Cation Entry and Exocytosis in Human Neutrophils: Lectins May Activate Multiple Subtypes of Cation Channels. *J. Leukocyte Biol.* **60**, 345-355.
71. Stewart, R., Allan, D.W. & McCraig, C.D. (1996). Lectins Implicate Specific Carbohydrate Domains in Electric Field Stimulated Nerve Growth and Guidance. *J. Neurobiol.* **30**, 425-437.
72. Rott, O., Charreire, J. & Cash, E. (1996). Influenza A Virus Hemagglutinin is a B-cell-superstimulatory Lectin. *Med. Microbiol. Immunol.* **184**, 185-193.
73. Singh, R.S., Tiwary, A.K. & Kennedy, J.F. (1999). Lectins: Sources, Activities, and Applications. *Crit. Rev. Biotech.* **19**, 145-178.
74. Blasioli, J. Paust, S. & Thomas, M.L. (1999). Definition of the Sites of Interaction between the Protein Tyrosine Phosphatase SHP-1 and CD22. *J. Biol. Chem.* **274**, 2303-2307.
75. Kemp, B. E., Rylatt, D. B., Bundesen, P. G., et al. (1988). Autologous red cell agglutination assay for HIV-1 antibodies: Simplified test with whole blood. *Science* **241**, 1352-1354.
76. Kawauchi, H., Hosono, M., Takayanagi, Y. & Nitta, K. (1993). Agglutinins from aquatic insects-tumor cell agglutination activity. *Experientia* **49**, 358-361.
77. Coloma, M. J. & Morrison, S. L. (1997). Design and production of novel tetravalent bispecific antibodies. *Nature Biotechnol.* **15**, 159-163.
78. Segal, D. M., Weiner, G. J. & Weiner, L. M. (1999). Bispecific antibodies in cancer therapy. *Curr. Opin. Immunol.* **11**, 558-562.
79. Furata, M., Uchikawa, M., Ueda, Y., et al. (1998). Construction of mono- and bivalent human single-chain Fv fragments against the D antigen in the Rh blood group: Multimerization effect on cell agglutination and application to blood typing. *Protein Eng.* **11**, 233-241.
80. Singh, R. S., Tiwary, A. K. & Kennedy, J. F. (1999). Lectins: Sources, activities, and applications. *Crit. Rev. Biotech.* **19**, 145-178.




81. Matsuya, Y. & Yamane, I. (1985). Cell hybridization and cell agglutination 1. Enhancement of cell hybridization by lectins. *J. Cell Sci.* **78**, 263-271.
82. Takamatsu, H., Kawajiri, H., Takahashi, Y., Manaf Ali, A. & Yoshimoto, T. (1999). Continuous antibody production by phytohemagglutinin-L-aggregated hybridoma cells. *J. Immunol. Methods* **223**, 165-170.
83. Crothers, D. M. & Metzger, H. (1972). The influence of polyvalency on the binding properties of antibodies. *Immunochem.* **9**, 341-357.
84. Mammen, M., Choi, S.-K. & Whitesides, G. M. (1998). Polyvalent interactions in biological systems: implications for design and use of multivalent ligands and inhibitors. *Angew. Chem. Int. Ed. Engl.* **37**, 2755-2794.
85. Kiessling, L. L. & Pohl, N. L. (1996). Strength in numbers: Non-natural polyvalent carbohydrate derivatives. *Chem. Biol.* **3**, 71-77.
86. Torti, M., Ramaschi, G., Sinigaglia, F. & Balduini, C. (1995). Dual mechanism of protein-tyrosine phosphorylation in concanavalin A-stimulated platelets. *J. Cell. Biochem.* **57**, 30-38.
87. Osawa, T. & Beppu, M. (1987). Cross-linked derivatives of concanavalin A. *Methods Enzymol.* **150**, 17-28.
88. Hudson, P. J. & Kortt, A. A. (1999). High avidity scFv multimers; diabodies and triabodies. *J. Immunol. Methods* **231**, 177-189.
89. Weis, W. I., Taylor, M. E. & Drickamer, K. (1998). The C-type lectin superfamily in the immune system. *Immunol. Rev.* **163**, 19-34.
90. Khan, M. I., Mandal, D. K. & Brewer, C. F. (1991). Interactions of concanavalin A with glycoproteins. A quantitative precipitation study of concanavalin A with the soybean agglutinin. *Carbohydr. Res.* **213**, 69-77.
91. Ballerstadt, R. & Schultz, J. S. (1997). Competitive-binding assay method based on fluorescence quenching of ligand held in close proximity by a multivalent receptor. *Anal. Chim. Acta* **345**, 203-212.
92. Burke, S. D., Zhao, Q., Schuster, M. C. & Kiessling, L. L. (2000). Synergistic formation of soluble lectin clusters by a templated multivalent saccharide ligand. *J. Am. Chem. Soc.* **122**, 4518-4519.

09615296-072004  
100E20-96251860


94. Gordon, E. J., Gestwicki, J. E., Strong, L. E. & Kiessling, L. L. (2000). Synthesis of end-labeled multivalent ligands for exploring cell-surface-receptor-ligand interactions. *Chem. Biol.* **7**, 9-16.
95. Gupta D, Dam TK, Oscarson S, Brewer CF.(1997) Thermodynamics of lectin-carbohydrate interactions. Binding of the core trimannoside of asparagine-linked carbohydrates and deoxy analogs to concanavalin A. *J Biol Chem.* **272**(10):6388-92.
96. Gestwicki, J.E., Strong, L. E. & Kiessling, L. L.(2000) Visualization of single multivalent receptor-ligand complexes by transmission electron microscopy. *Angew. Chem. Int. Ed.* **39**, No. 24, 4567-4570.
97. Parkin, S., Rupp, B., Hope, H. (1996) Atomic resolution structure of concanavalin A. *Acta Crystallogr. Sect. D* **52**:1161-1168.
98. Hunig, S., Muller, H.R., Their, W. *Angew. Chem. Int. Ed. Engl.* (1965) **4**:271-382.
99. Hart, D.J.; Hong, W.-P.; Hsu, L-Y. *J. Org. Chem.* (1987) **52**: 4665-4673.
100. Johnson, R.A., Sharpless, K.B. (1993) In *Catalytic Asymmetric Synthesis*, Ojima, I. (ed.)VCH: New York .
101. Walsh, P.J., Bennani, Y.L., Sharpless, K.B. (1993) *Tetrahedron Lett.* **34**:5545-5548.
102. Kiss et al. (1997) *Gut* **40**:253.

All of the references cited herein are incorporated by reference herein in their entirety and to the extent that they are not inconsistent with the disclosures herein. The cited references are incorporated by reference herein in particular for any description regarding the synthesis of multivalent ligands and particularly synthesis by ROMP methods and for description regarding the selection of signal groups, particularly chemoattractants or epitopes, for a given application.


# SCHEME 1: MULTIVALENT ROMP POLYMERS WITH SACCHARIDE SRE




SCHEME 2: ROMP POLYMERS FOR NEUTROPHIL CHEMOTAXIS




SCHEME 3: ROMP → PEPTIDE LINKERS




SCHEME 4: EXAMPLE LIGANDS FOR BINDING TO B CELLS



SCHEME 5:  
EXEMPLARY RE AND SRE FOR  
MULTIVALENT LIGANDS THAT BIND TO B CELLS



50




51

09815296-073001



SCHEME 6: METHODS FOR MODIFYING POLYMER BACKBONES



SCHEME 7: SYNTHESIS OF A CAPPING AGENT

